

IL 99.417



STATE OF ISRAEL

09 / 744804

REC'D 01 SEP 1999

WIPO PCT

This is to certify that
annexed hereto is a true
copy of the documents as
originally deposited with
the patent application
particulars of which are
specified on the first page
of the annex.

זאת לתעודה כי
רצופים בזה העתקים
נכונים של המסמכים
שהופקרו לכתחילה
עם הבקשה לפטנט
לפי הפרטים הרשומים
בעמוד הראשון של
הנספח.

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

01-08-1999

This היום

מ.ל.ס. 1000
מ.ל.ס. 1000

רשם הפטנטים
Registrar of Patents

נתאשר

PATENT LAW, 5727 - 1967

PATENT LAW, 5727 - 1967

ב.ק.ש.ה ל פ ט נ ט

Application For Patent

אני, (שם המבקש, מענו ולגבי גוף מאוגד - מקום התאגדותו)

1 (Name and address of applicant, and in case of body corporate - place of incorporation)

Yeda Research And Development Company Ltd.
P.O.Box 95
Rehovot 76100

ידע חברה למחקר ופתוח בע"מ

95. ת.ת

76100 רחובות

שם המצאת.....Assignment.....בעל אמצאה מכה העברה
of an invention the title of which is Owner, by virtue of

פפטידים הקשורים לגידולים סרטניים ושימוש בהם כחיסון נגד גידולים סרטניים

(בעברית)

(Hebrew)


TUMOR ASSOCIATED ANTIGEN PEPTIDES AND US OF SAME AS ANTI-TUMOR VACCINES

(באנגלית)

(English)

hereby apply for a patent to be granted to me in respect thereof

מבקש בזאת כי ינתן לי עליה פטנט

• בקשת חלוקה - Application of Division		• בקשת פטנט מוסף - Application for Patent Addition		• דרישת דין קדימה Priority Claim	
מבקשת פטנט from Application No. מס' dated מיום		לבקשה/לפטנט to Patent/Appl. No. מס' dated מיום		מספר/סימן Number/Mark	תאריך Date
יפוי כח : כללי P.O.A.: general/individual-attached/to be filed later- הוגש בענין 104513 filed in case		המען למסירת מכתבים בישראל Address for service in Israel DR. MARK FRIEDMAN LTD. BEIT SAMUELOFF 7 HAOMANIM STREET 67897 TEL AVIV			
חתימת המבקש Signature of Applicant 		1998 שנת Jul of the year		30 בחודש of This	לשימוש הלשכה For Office Use

Ref.: 34/30

פפטידים הקשורים לגידולים סרטניים ושימוש בהם כחיסון נגד גידולים סרטניים

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS
ANTI-TUMOR VACCINES

TUMOR ASSOCIATED ANTIGEN PEPTIDES AND USE OF SAME AS ANTI-TUMOR VACCINES

FIELD AND BACKGROUND OF THE INVENTION

5 The present invention relates to tumor associated antigen (TAA) peptides and to the use of same, of polynucleotides encoding same and of cells presenting same as anti-tumor vaccines. More particularly, the present invention relates to a tumor associated antigen peptides derived from Uroplakin II and Ia, Prostate specific antigen (PSA), Prostate acid phosphatase (PAP) and Prostate specific
10 membrane antigen (PSMA), BA46 (Lactadherin), Mucin (MUC-1), and the use of same as anti-tumor vaccines to prevent or cure bladder, prostate, breast or other cancers, carcinomas in particular.

 Local therapy such as surgical excision or ablation by radiation is a mainstay for the treatment of primary cancer and is curative for a percentage of
15 patients. However, many malignancies will recur locally or at a distant site. Thus the prevention or cure of metastases remains a major focus in clinical oncology (1). Although early detection followed by surgery provides good prognosis for a number of major cancer types, a large fraction of patients would need adjuvant therapy. Part of these patients will, with time, succumb to
20 metastasis (2-4). Alternative approaches based on gene therapy and immunotherapy have focused attention in the last years. One such approach is specific active immunotherapy (SAI, 5). The objective of SAI is to stimulate a tumor specific cytotoxic T lymphocytes (CTL) immune response that is capable of eliminating residual metastatic disease and induce a state of immunity to
25 protect the patients from recurrent disease. The underlying assumption of SAI is that tumor cells express tumor antigens which are sufficiently distinct in structure or context to induce an effective CTL response (6). Although the validity of these assumptions was questioned, a number of studies in the last decade have demonstrated the rational of ASI. In a landmark study, van Pel and Boon have
30 shown that tumor associated antigens (TAAs) can be isolated and defined (7). Importantly, *ex-vivo* manipulations of "non-immunogenic" animal tumor cells can be used to elicit effective immune responses which will also recognize parental "non-immunogenic" tumor cells (8). Studies employing rodent tumor models with little intrinsic immunogenicity have shown that genetically modified
35 tumor cells vaccines transduced to express MHC class I, cytokines such as IL-1, IL-2, IL-4, IL-6, IL-7, IL-12, γ IFN or GM-CSF or costimulatory molecules such as B7-1 or B7-2 were capable of preventing or causing regression of tumors or metastases (reviewed in 9-12). Although gene modified tumor vaccine (GMTV) clinical trials, with improved retroviral vectors or other transfer methodologies

are currently tested, it becomes clear that GMTV using autologous tumor cells might be limited by its complexity, high cost and ineffective gene transfer methodologies (13). One alternative approach would be vaccination with tumor associated antigens (TAAs) presented in an effective way to the patient's immune system, to induce antigen specific CTL (14).

Cytotoxic T lymphocytes (CTL), directed against peptides presented by MHC class I molecules, constitute powerful effectors of the immune system against tumors or infectious agents (15). These peptides are usually 8-10 amino acid long with 2-3 primary anchor residues that interact with the MHC class I molecules and 2-3 amino acid residues which bind to the T cell receptor (16-18). Several methods have been employed to identify CTL epitopes. If the amino acid sequence of a protein antigen is known, like in the case of viral proteins, oncogenes, suppressor genes or growth factor receptors, overlapping peptides of 8-10 amino acids in length can be synthesized and screened as CTL targets (19, 20). CTL epitopes may also be identified subsequent to the search for MHC binding motifs in known proteins (21-24). If the tumor antigen is not known, isolation of the TAA peptides from total acid extract or from MHC class I molecules followed by HPLC fractionation steps and Edman sequencing (25, 26) or mass spectrometry (27) provided a direct way of identifying CTL epitopes. Recently, a synthetic combinatorial library approach, in which defined amino acids in two MHC anchor positions are fixed and all other positions are subgrouped for CTL screening has led to the description of novel EL4 TAA peptide mimotopes (28).

The most fruitful method, so far, designed by T. Boon and his colleagues is the genetic approach in which cDNA expression libraries are pool transfected into COS7 cells with the appropriate HLA and screened by CTL lines. This approach led to the discovery of several human melanoma and mouse mastocytoma antigens recognized by specific CTL (29-34). The first report of a phase I clinical trial with the synthetic MAGE3 melanoma peptide, restricted by HLA-A1, showed regression of cutaneous, subcutaneous and lung metastases in 3/6 patients (35). Recently, two reports of clinical trials have shown that treatment of patients with a melanoma gp100 TAA peptide together with IL-2 resulted in significant tumor regression in 13/31 (42 %) patients and that vaccination with defined peptides or total peptide extracts on autologous Dendritic cells (DC) resulted in complete or partial cures (36, 37). Regression of lung carcinoma established metastases or small established tumors was demonstrated by us in a murine model by peptide vaccination (38, 39). These observations suggest that TAA peptide vaccines may constitute a reasonable

therapeutic modality in advanced cancer. In studies with murine tumors, CTL are induced *in vivo* by immunization with irradiated tumor cells, often gene modified by MHC class I; cytokine or costimulatory molecules like B7-1 or B7-2 genes (26, 28, 40-42). In melanomas, CTL lines were mostly induced from peripheral
 5 blood mononuclear cells (PBMC) of patients or from tumor infiltrated lymphocytes (TiL, 31, 43-47). Yet, most metastatic tumors are non-immunogenic tumors and it is extremely difficult to derive CTL lines or clones from TiL or patient's PBL. Moreover, *in vitro* propagated CTL clones do not always represent dominant anti-tumor specificities but rather sporadic clones
 10 surviving culture conditions. Lately, a number of studies have compared the CTL repertoire of viral or other defined peptides, restricted by HLA-A2.1 in human PBL from HLA-A2.1 expressing patients to CTL induced in HLA-A2.1 transgenic mice. Good concordance between human HLA-A2.1 and murine transgenic HLA-A2.1 CTL repertoire was found, confirming the potential of such
 15 transgenics in identification of human CTL epitopes (48-52). Although vaccination with defined peptides of HLA transgenic mice shows an overlapping repertoire to human CTL, vaccination of such mice with multi-epitope proteins shows that murine H-2 restricted responses are dominant and obliterate, as a rule, cytolytic responses with direct recognition of human HLA (53). Thus, by
 20 combining classical HLA class I transgenesis with selective destruction of murine H-2, it is possible to derive useful mouse strains for the study of HLA class I restricted responses. While reducing the present invention to practice, we utilized such mice for preparation of anti-tumor CTL as a tool for TAA purification and as a model system to assess the immunogenicity of peptides.

25 Murine H-2 knockout mice transgenic for a single human HLA seem to be a suitable model for induction of anti-tumor CTL. Classical $\beta 2$ microglobulin knockout mice ($\beta 2m^{-/-}$) do not express H-2K^b or other non-classical class I molecules, yet they express low levels of H-2D^b heavy chain in the absence of $\beta 2m$. To derive fully H-2 knockout mice, Prof. F. Lemonnier (Pasteur Institute,
 30 Paris), prepared H-2D^b $^{-/-}$ mice. These mice were crossed with $\beta 2m^{-/-}$ mice and bred to derive homozygous $\beta 2m^{-/-}$, D^b $^{-/-}$ mice that do not express any H-2 class I. These mice are practically depleted of CD8⁺ splenocytes, as well as other CD8⁺ cells. To reconstitute in these mice expression of a stable HLA-A2.1, expression of $\beta 2m$ is necessary. A construct containing a leader sequence, $\alpha 1$
 35 and $\alpha 2$ domains of HLA-A2.1 and $\alpha 3$, transmembrane and cytoplasmic domains of H-2D^b fused to human $\beta 2m$ (HhD) was prepared. The exchange of the $\alpha 3$ human domain by a murine domain in HhD is thought to improve the interaction of the class I molecule with CD8 molecules of the murine CTL (54). This HhD

construct was transfected into RMA and RMA-S cells and shown to bind HLA-A2.1 restricted peptides. The HhD construct was used to produce transgenic mice in C57BL/6 recipients and positive founder mice were bred to the $\beta_2m^{-/-}$, $Db^{-/-}$ mice (55).

5 The $\beta_2m^{-/-}$, $Db^{-/-}$, $HhD^{-/+}$ heterozygous mice show reconstitution of $CD8^{+}$ cells in the periphery relative to $\beta_2m^{-/-}$ $Db^{-/-}$ mice. Moreover, preliminary data from Prof. Lemonnier's lab showed that CTL induced in HhD mice against influenza NP are directed to the same HLA-A2 dominant epitope as in the human repertoire. Homozygous HhD mice were derived and a colony was
10 established in the Weizmann Institute of Science, Israel.

15

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a vaccine composition comprising at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP),
20 Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

According to another aspect of the present invention there is provided a method of vaccination for prevention or cure of cancer comprising the steps of administering a composition including at least one tumor associated antigen
25 peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

According to yet another aspect of the present invention there is provided
30 a DNA vaccine composition comprising at least one polynucleotide encoding a tumor associated antigen peptide derived from a sequence encoding a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1). The at least one polynucleotide
35 can be a part of a longer polynucleotide designed to encode a fused protein product from which the tumor associated antigen peptide is cleavable by a protease.

According to still another aspect of the present invention there is provided a cellular vaccine composition comprising an antigen presenting cell presenting at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1). The antigen presenting cell can, for example, be a dendritic cell, a macrophage, a B cell and a fibroblast. Presenting the at least one tumor associated antigen peptide can be effected by a method selected from the group consisting of (a) transducing the antigen presenting cell with at least one polynucleotide encoding the at least one tumor associated antigen peptide; (b) loading the antigen presenting cell with at least one polynucleotide encoding the at least one tumor associated antigen peptide; (c) loading the antigen presenting cell with the at least one tumor associated antigen peptide; (d) loading the antigen presenting cell with at least one longer polypeptide including the at least one tumor associated antigen peptide.

According to further features in preferred embodiments of the invention described below, the Uroplakin is selected from the group consisting of Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.

According to still further features in the described preferred embodiments the one or more tumor associated antigen peptides derived from Mucin are from a non tandem repeat array of the Mucin.

According to still further features in the described preferred embodiments the one or more tumor associated antigen peptides derived from Mucin are from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of Mucin.

According to still further features in the described preferred embodiments the Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1) are each independently of a mammal.

According to still further features in the described preferred embodiments the mammal is selected from the group consisting of humanoid and rodent.

According to still further features in the described preferred embodiments each of the one or more tumor associated antigen peptide includes a number of residues including 2-3 residues which form a primary anchor for interaction with MHC class I molecules and 2-3 residues which bind to T cell receptors.

According to still further features in the described preferred embodiments the number is selected from the group consisting of 7, 8, 9, 10 and 11.

7

According to still further features in the described preferred embodiments the composition further comprising a carrier, wherein the tumor associated antigen peptide(s) are presented in context of the carrier.

5 According to still further features in the described preferred embodiments the carrier is selected from the group consisting of a proteinaceous carrier to which said at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.

10 According to still further features in the described preferred embodiments at least one of the tumor associated antigen peptides includes at least one non-natural modification rendering peptides more stable in a body.

According to still further features in the described preferred embodiments at least one of the tumor associated antigen peptides includes at least one non-natural modification rendering peptides more immunogenic.

15 According to still further features in the described preferred embodiments the non-natural modification rendering peptides more immunogenic and/or more stable in the body is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

20 According to still further features in the described preferred embodiments the one or more tumor associated antigen peptides are selected from SEQ ID NOs: 1-49 and effective homologues and analogs thereof.

According to still further features in the described preferred embodiments the composition is effective in prevention or cure of cancer or cancer metastases.

25 According to still further features in the described preferred embodiments the cancer is a carcinoma.

According to still further features in the described preferred embodiments the cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, melanoma, colon, stomach and head and neck cancers.

30 The present invention successfully addresses the shortcomings of the presently known configurations by providing novel tumor associated antigen peptides effective in eliciting CTL response which may therefore be effective therapeutic agents to combat cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention herein described, by way of example only, with reference to the accompanying drawings, wherein:

FIG. 1 demonstrates stabilization of cell surface HhD on RMA-S cells by human Uroplakin II derived peptides. Uroplakin II derived synthetic peptides (HURO1-7) or a peptide from the melanoma associated TAA Tyrosinase (Tyr) which served as a control were loaded at various concentrations (3-300 μ M) on the RMA-HhD cells as described. Indirect fluorescence activated cell sorter (FACS) analyses were performed by incubating 5×10^5 loaded cells with the anti-HLA-A2 monoclonal antibodies BB7.2 for 30 minutes at 4 °C. After washing the cells in PBS-0.5 %, BSA-0.1 %, sodium azide, the second antibody, goat anti mouse-FITC was applied for 30 minutes at 4 °C. Following another wash the fluorescence was recorded on a FACSScan (BD). Mean fluorescence at 300 μ M is shown.

FIG. 2 demonstrates lytic activity of human peripheral blood lymphocytes (PBL) against Uroplakin II loaded target cells. PBL from HLA-A2 positive patients were resensitized with tumor extracted peptide preparations from transitional cell carcinoma (TCC) specimens as described in Materials and Methods. Resensitized lymphocytes from 4 TCC patients treated with Bacillus Calmette Guerein (BCG), 2 non-treated TCC patients, and 1 prostate cancer patient (control), were admixed at different ratios with labeled T2 cells, loaded with TCC extracted peptides (TCC pool) normal bladder mucosa extracted peptides (bladder), synthetic Uroplakin II peptides (HURO1-7) or the melanoma peptide from Tyrosinase (Tyr). Labeled target cells, J82, a HLA-A2 expressing TCC line, SUP, a low class I expressor and k562, a NK sensitive line were also tested. Percent specific lysis, at effector: target of 100:1 is presented.

FIG. 3 demonstrates lytic activity of human peripheral blood lymphocytes against peptides extracted from individual TCC tumors. PBL were resensitized as in Figure 2. Peptides extracted from 9 individual tumors (TCC1-9), that were pooled for resensitization, were loaded on T2 cells. Labeled loaded targets were tested for lysis by sensitized PBL as described in Figure 2.

FIG. 4 demonstrates lysis of human Uroplakin II and murine Uroplakin II peptide loaded target cells by CTL induced again human Uroplakin II derived peptides in HhD mice. Mice were immunized as described, by RMA-S-HhD-B7.1 cells, loaded with human Uroplakin II derived synthetic peptides 1-7 as described in Materials and Methods. Loading was performed with individual peptides and cells were pooled for immunization. Resensitized CTL (see Materials and Methods) were tested for lysis of RMA-S-HhD targets loaded with

human or murine Uroplakin II peptides (HURO 1-7, MURO1, 3, 4, 6) with TCC or normal bladder extracted peptides (RMAS TCC, RMAS bladder), with the tyrosinase peptide (RMAS Tyr) with peptides derived from breast tumors (RMAS Br.) or with the cell lines J82 or SUP. E:T ratios of 100:1, 50:1, 25:1 and
 5 12.5:1 were tested and the results are presented in lytic units LU30 was calculated by linear regression analysis of percentage lysis vs. the log (ln) of effector cell number and expressed as the number of LU per 10^6 effector cells. The coefficient was 0.85 - 0.95 for all groups.

FIG. 5 demonstrates lysis of human Uroplakin II and Ia and murine
 10 Uroplakin II derived peptide loaded target cells by CTL induced against TCC extracted peptides in HhD mice. HhD mice were immunized by RMA-S-HhD-B7.1 cells loaded with peptides extracted from 9 TCC samples and pooled. CTL were prepared as described and tested for lysis of labeled RMA-S-HhD target cells loaded with synthetic, tissue derived or control peptides and cell lines as in
 15 Figure 4 and synthetic peptides derived from Uroplakin Ia (HURO 11-18). The results are presented in lytic units as described in Figure 4.

FIG. 6 demonstrates stabilization of cell surface HhD on RMA-S cells by human Prostate specific antigen (PSA) and Prostate acid phosphatase (PAP) derived peptides. PSA and PAP derived peptides were loaded at various
 20 concentrations (1 μ M - 1 mM) on RMA-S-HhD cells. FACS analysis was performed as described in Figure 1, wherein Tyr served as a control. Mean fluorescence at 1 mM is presented.

FIG. 7 demonstrates stabilization of cell surface HhD on RMA-S cells by human Prostate specific membrane antigen (PSMA) derived peptides. PSMA
 25 derived peptides (1 μ M - 1 mM) were loaded and cells were monitored as described in Figure 6.

FIG. 8 demonstrates immunogenicity and antigenicity of PAP derived peptides in HhD mice. HhD mice were immunized as described, by RMA-S-HhD-B7.1 cells loaded individually with each PAP peptide and pooled before
 30 vaccination. CTL were tested against RMA-HhD cells loaded with synthetic peptides (PAP 1-4), prostate carcinoma extracted peptides (RMAS, CAP), hyperplastic (normal) prostate extracted peptides (RMAS HP), breast carcinoma derived peptides (RMAS Br, control) and tyrosinase (RMAS Tyr, control). The prostate carcinoma cell line DU145 (DU) and its HhD transfectant (DU HhD)
 35 were also tested as targets. The results are presented in lytic units as in Figure 4.

FIG. 9 demonstrates immunogenicity and antigenicity of PSA derived peptides in HhD mice. Procedures were performed and results are presented as in Figure 8.

FIG. 10 demonstrates immunogenicity and antigenicity of PSMA derived peptides in HhD mice. Procedures were performed and results are presented as in Figure 8.

FIG. 11 demonstrates lysis of PSA, PAP and PSMA derived peptide loaded RMA-S-HhD cells by CTL induced against DU145-HhD cells in HhD mice. Mice were immunized with irradiated DU145-HhD cells and CTL assays were performed as described in Materials and Methods. Labeled RMA-S-HhD target cells were loaded with synthetic PSA (1-5), PAP (1-4), PSMA (1-6) or tyrosinase (RMAS Tyr) peptides, with prostate carcinoma (RMAS CAP), breast carcinoma (RMAS Br.) or hyperplastic prostate (RMAS HP) extracted peptides. DU145 (DU) and DU145-HhD also served as targets. The results are presented in lytic units as in Figure 4.

FIG. 12 demonstrates lysis of PSA, PAP and PSMA derived peptide loaded RMA-S-HhD by CTL induced against prostate carcinoma extracted peptides in HhD mice. HhD mice were immunized with RMA-S-HhD-B7.1 cells loaded with peptides extracted from prostate carcinoma samples. Experimental details are equivalent to the details in Figure 11.

FIG. 13 demonstrates stabilization of cell surface HhD on RMA-S- cells by human BA46 (Lactadherin) derived peptides. RMA-S-HhD cells were incubated with 1-100 μ M synthetic peptides and monitored as described in Figure 1.

FIG. 14 demonstrates immunogenicity of BA46 peptides in HhD mice. Mice were immunized as described in Materials and Methods, with RMA-S-HhD-B7.1 cells loaded with synthetic peptides. CTL assays were performed on RMA-S-HhD target cells loaded with homologous peptides.

FIG. 15 demonstrates lysis of BA46 peptide loaded targets by CTL induced against breast carcinoma extracted peptides in HhD mice. HhD mice were immunized with RMA-S-HhD-B7.1 cells loaded with peptides extracted from breast carcinoma samples. Experimental details are as in Figure 11. The data is presented as percent specific lysis at E:T of 100:1.

FIG. 16 demonstrates differential lysis of tumor extracted peptide loaded target cells by CTL induced against tumor peptides or BA46 derived peptides in HhD mice. Mice were immunized by RMA-S-HhD-B7.1 cells loaded with synthetic BA46 derived peptides or breast carcinoma derived peptides as described in Materials and Methods. CTL were tested against RMA-S-HhD cells loaded with breast carcinoma extracted peptides or normal breast tissue extracted peptides. Percent specific lysis at E:T of 100:1 is presented.

FIG. 17 demonstrates HLA-A2.1 (HhD) restricted lysis of a breast carcinoma cell line by CTL induced against BA46 derived peptides or tumor extracted peptides. HhD mice were immunized as in Figure 16 and CTL were tested against the HLA negative MDA-MB-157 and their HhD transfectants. The data is presented in percent specific lysis at a E:T of 100:1.

FIG. 18 demonstrates stabilization of cell surface HhD on RMA-S- cells by Mucin (MUC-1/A7, E6, D6) derived peptides. Details are as described in Figure 1.

FIG. 19 demonstrates immunogenicity of MUC-1 derived peptides. HhD mice were immunized and CTL tested as in Figure 14.

FIG. 20 demonstrates HLA-A2.1 (HhD) restricted lysis of a breast carcinoma cell line by CTL induced against MUC-1 derived peptides. Details as described in Figure 17.

FIG. 21 demonstrates lysis of MUC-1 loaded target cells by CTL induced against breast carcinoma extracted peptides in HhD mice. Details are as described in Figure 15. E:T is 50:1.

FIG. 22 demonstrates differential lysis of tumor extracted peptide loaded target cells by CTL induced against MUC-1 derived peptides in HhD mice. Details are as described in Figure 16.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of a vaccine composition which can be used to prevent or cure cancer, both primary tumors and metastases. Specifically, the present invention can be used to provide vaccines which include novel and potent tumor associated antigen peptides derived from Uroplakins, Prostate specific antigen, Prostate acid phosphatase, Prostate specific membrane antigen, Lactadherin and Mucin, which can be used as anti-tumor vaccines to prevent or cure, for example, bladder, prostate or breast cancers, carcinomas in particular, or any other tumor expressing the above listed proteins.

The principles and operation of the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other

embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

5 A specific CTL response was found in patients with bladder transitional cell carcinoma (TCC) against the tumor extract and against peptides homologue to Uroplakin II sequence, which was significantly augmented after BCG intravesical therapy. All Uroplakin II peptides employed were good immunizers in HhD mice and CTL from these mice induced intense lysis of targets loaded with Uroplakin II homologue peptides, TCC peptide extract and of J82 9HLA-A2
10 expressing cell line. CTL from HhD mice immunized with TCC peptide extract showed cross reactivity and induced significant lysis of targets loaded with Uroplakin II, or Uroplakin Ia homologue peptides, with TCC peptide extract and of J82 cells. For all HhD mice, there was reduced lysis of targets loaded with normal bladder mucosa peptide extract and of TCC SUP cells (minimal HLA-A2
15 expression). Also there was no non-specific lysis. Also, a cross reactivity of CTL induced by human Uroplakin II peptides against murine homologue peptides was found. Despite this finding there was no damage, or inflammatory infiltrate in the internal organs, including the urinary bladder on careful histological examination. These data indicate that Uroplakin homologue peptides
20 constitute specific CTL epitopes and may be considered for immunotherapeutic vaccines alone, or in combination with intravesical BCG instillations in patients prone to recurrence and progression of bladder TCC.

By using HhD mice, HLA-A2 transgenic and which do not express murine MHC class I, we found 3 Prostate specific antigen (PSA), 4 Prostate specific
25 membrane antigen (PSMA) and 4 Prostate acid phosphatase (PAP) homologue peptides to be immunogenic. The 3 PSA homologue peptides and 2 of the 4 PSMA homologue peptides were not evaluated previously. The 2 PSMA homologue peptides, which were evaluated previously and were found to be CTL epitopes in CAP patients were also found by us to be immunogenic in HhD mice.
30 PAP homologue peptides were not evaluated previously and we found that all 4 HLA-A2 binding peptides were very immunogenic in HhD mice and induced also, specific lysis of DU145-HhD cells. Moreover, we found that CTL derived from HhD mice immunized with CAP peptide extract, or with DU145-HhD cells induced specific lysis of targets loaded with CAP peptide extract and PAP
35 homologue peptides. There was also, cross reactivity between CTL derived from CAP immunized HhD mice and PSA, or PSMA homologue peptide loaded targets. Low cross reactivity was found for CTL derived from DU145-HhD immunized HhD mice, probably due to low expression of these proteins by

DU145 cells. There was no damage, or inflammatory infiltrate in the internal organs of immunized or control HhD mice on careful histological examination. These data indicate that PSA, PSMA and especially PAP homologue peptides constitute specific CTL epitopes and may be considered for immunotherapy of CAP patients. We intend to evaluate the value of adjuvant immunization with PSA, PSMA and PAP homologue peptides in HLA-A2 patients after radical prostatectomy, and in patients with metastatic prostate cancer.

We further found a specific CTL response in HhD mice against breast tumor extract and against novel peptides complementary to Lactadherin (BA46) sequences. These data indicate that BA46 peptides constitute specific CTL epitopes enriched in breast carcinoma and may be considered for immunotherapeutic vaccines.

Yet we further found that Mucin (MUC-1) derived peptides are potential TAA peptides that can be used in anti-tumor vaccine preparations. CTL induced by these peptides lyse better tumor extract loaded targets than normal tissue extract loaded targets. The lysis is HLA-A2 restricted and breast specific. Three novel peptides, from non tandem repeat array (TRA) domains were shown to constitute potential CTL epitopes.

Thus, in accordance with one aspect of the teachings of the present invention there is provided a vaccine composition which includes at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

According to another aspect of the present invention there is provided a method of vaccination for prevention or cure of cancer. The method is effected by administering a composition including at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

Immunotherapy by *in vivo* DNA transfer of DNA coding for TAA is based on the rationale of quality or quantity increased peptide presentation leading to activation of an immune response against these peptides. Gene or DNA vaccination results in the intracellular processing and presentation of immunogenic peptides (99). Initial reports on DNA vaccination showed that "naked" DNA injected into the muscle tissue of a mouse is expressed efficiently (100). Embryonically expressed TAA such as CEA was tested (101).

Immunization of mice with CEA expressing plasmid DNA was indeed found to protect 100 % of these mice against a challenge with CEA-expressing colon carcinoma cells (102). Both cellular and humoral responses have been reported after DNA vaccination in mice. In other studies a MUC-1 tandem repeat array was used for DNA vaccination of mice and 30% of these mice were protected from a tumor challenge with MUC-1 transfected murine tumor cells (103). DNA vaccination may also be used to elicit immune responses against predefined peptide epitopes, several groups now exploit the string-bead approach to link multiple different CTL or helper epitopes together on the DNA level (104, 105). In some cases the string-bead of peptide coding DNA is built into a vaccinia virus as a delivery vehicle. Recently, it was shown that such a vaccinia virus recombinant poly-epitope vaccine was able to protect mice against several virus infections and a tumor challenge (106, 107). The authors show that all 10 minimal peptide epitopes encoded by the string-bead are expressed and recognized by the appropriate T cell clones (108). RNA was also shown to confer anti-tumor immunity. Vaccination with RNA to ovalbumin induced CTL in mice (109). In conclusion, multiple studies have shown the efficacy of DNA vaccines in anti-viral and anti-tumor immunity.

Thus, according to yet another aspect of the present invention there is provided a DNA vaccine composition which includes at least one polynucleotide encoding a tumor associated antigen peptide derived from a sequence encoding a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1). As further detailed hereinunder, the at least one polynucleotide can be a part of a longer polynucleotide designed to encode a fused protein product from which the tumor associated antigen peptide is cleavable by a protease.

The polynucleotide is preferably DNA in a form of, or contained in, for example, naked DNA, plasmid, retroviral vector, adenoviral vector, vaccinia viral vector, herpes viral vector, lenti virus vector, EBV vector, CMV vector, polio virus vector, sindbis viral vector, semliki forest virus vector, parvo virus vector, adeno-associated virus vector, virus like particle (VLP) vector. Alternatively, the polynucleotide can be in the form of RNA. The polynucleotide can also be delivered in a non-viral delivery system, such as, for example, but not limited to, in liposomes, in complex with cationic reagents, or with a polycation, such as poly-lysine. The polynucleotide can also be delivered by mechanical means, such as, but not limited to, a gene-gun, by electrical means, or in bacterial vectors like BCG.

There is increasing evidence that peptide vaccination may be much more effective when the peptides are introduced together with an antigen presenting cell (APC, 110, 111). In previous studies of a murine lung carcinoma we have shown that vaccination with a defined TAA peptide (MUT-1) loaded on APC
 5 result in long term survival of mice bearing lung metastasis (112, 113). The most common cells used to load antigens are bone marrow and peripheral blood derived dendritic cells (DC), as these cells express costimulatory molecules that help activation of CTL. Preliminary clinical trials have been performed. In one trial HLA-A1 melanoma patients have been treated with autologous DC loaded
 10 with a MAGE-1 peptide. CTL activity was increased in tumor infiltrated lymphocytes (114). In another study, five patients with advanced pancreatic carcinoma were treated with a K-ras derived peptide loaded on DC. As a mutation of K-ras at codon 12 is frequently found in pancreatic carcinoma, three differently mutated peptides, 12-Asp, 12-Arg and 12-Val (non mutated sequence
 15 is 12-Gly) were used for vaccination, matched to the mutation in the patient's tumor. Two of the patients showed a specific CTL response and prolonged survival (115). A phase I clinical trial in 51 prostate cancer patients compared a soluble peptide to a DC based peptide in HLA-A2 patients. The peptide was derived from PSMA (SEQ ID NO:29). Only 7 patients that received DC based
 20 vaccines with this peptide responded by decreased levels of serum PSA (116). In animal studies a number of groups showed that macrophages loaded with peptides constitute efficient vaccines, yet the number of cells used for vaccination is 10 fold higher than equivalent DC vaccines. Recently we tested in a murine lung carcinoma model the efficacy of syngeneic fibroblasts, treated with a
 25 proteasome inhibitor to decrease levels of endogenous peptides and loaded with synthetic MUT peptides as vaccines. We found effective protection against metastatic spread of lung carcinoma.

Thus, according to still another aspect of the present invention there is provided a cellular vaccine composition which includes an antigen presenting cell
 30 presenting at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1). As further detailed hereinunder, the antigen presenting cell can, for example, be a dendritic cell, a
 35 macrophage, a B cell and a fibroblast. Presenting the at least one tumor associated antigen peptide can be effected by a method selected from the group consisting of (a) transducing the antigen presenting cell with at least one polynucleotide (e.g., DNA) encoding the at least one tumor associated antigen

peptide; (b) loading the antigen presenting cell with at least one polynucleotide (e.g., RNA) encoding the at least one tumor associated antigen peptide; (c) loading the antigen presenting cell with the at least one tumor associated antigen peptide (e.g., synthetic); (d) loading the antigen presenting cell with at least one
 5 longer polypeptide (e.g., purified) including the at least one tumor associated antigen peptide. Loading can be external or internal. The polynucleotide, peptide or longer polypeptide can be fused to internalizing sequences, antennapedia sequences or toxoid sequences or to helper sequences, such as, but not limited to, heat shock protein sequences.

10 As used herein in the specification and in the claims section below the phrase "tumor associated antigen" also refers to tumor specific antigen.

As used herein in the specification and in the claims section below the term "peptide" refers to native peptides (either degradation products or synthetically synthesized peptides) and further to peptidomimetics, such as
 15 peptoids and semipeptoids which are peptide analogs, which may have, for example, modifications rendering the peptides more stable while in a body, or more immunogenic. Such modifications include, but are not limited to, cyclization, N terminus modification, C terminus modification, peptide bond modification, including, but not limited to, CH₂-NH, CH₂-S, CH₂-S=O, O=C-
 20 NH, CH₂-O, CH₂-CH₂, S=C-NH, CH=CH or CF=CH, backbone modification and residue modification. Methods for preparing peptidomimetic compounds are well known in the art and are specified in Quantitative Drug Design, C.A. Ramsden Gd., Chapter 17.2, F. Choplin Pergamon Press (1992), which is incorporated by reference as if fully set forth herein. Further detail in this respect
 25 are provided hereinunder.

As used herein in the specification and in the claims section below the term "amino acid" is understood to include the 20 naturally occurring amino acids; those amino acids often modified post-translationally *in vivo*, including for example hydroxyproline, phosphoserine and phosphothreonine; and other
 30 unusual amino acids including, but not limited to, 2-amino adipic acid, hydroxylysine, isodesmosine, nor-valine, nor-leucine and ornithine. Furthermore, the term "amino acid" includes both D- and L-amino acids. Further elaboration of the possible amino acids usable according to the present invention and examples of non-natural amino acids useful in MHC-1 recognizable peptide
 35 antigens are given hereinunder.

Thus, assume the following positions (P1-P9) in a 9-mer peptide:

P1-P2-P3-P4-P5-P6-P7-P8-P9

Preferences of amino acids at positions P2 and P6 (anchor residues) and at positions P1 and P3 (secondary or auxiliary anchor residues) are given in Table V of reference 117.

Hydrophilic aliphatic natural amino acids at P2 and P9 can be substituted by synthetic amino acids, preferably Nleu, Nval and/or α -aminobutyric acid. P9 can be also substituted by aliphatic amino acids of the general formula $-\text{HN}(\text{CH}_2)_n\text{COOH}$, wherein $n = 3-5$, as well as by branched derivatives thereof, such as, but not limited to,



wherein R is, for example, methyl, ethyl or propyl, located at any one or more of the n carbons.

Each one, or more, of the amino acids at positions P1-P9 can include a D-isomer thereof. D isomers, however, are more accepted at positions P4-P8. In fact, the latter sites can accommodate most, if not all, amino acid residues, natural as well as non-natural, aliphatic and aromatic.

The amino terminal residue (position P1) can be substituted by positively charged aliphatic carboxylic acids, such as, but not limited to, $\text{H}_2\text{N}(\text{CH}_2)_n\text{COOH}$, wherein $n = 2-4$ and $\text{H}_2\text{N}-\text{C}(\text{NH})-\text{NH}(\text{CH}_2)_n\text{COOH}$, wherein $n = 2-3$, as well as by hydroxy Lysine, N-methyl Lysine or ornithine (Orn). Additionally, the amino terminal residue can be substituted by enlarged aromatic residues, such as, but not limited to, $\text{H}_2\text{N}-(\text{C}_6\text{H}_6)-\text{CH}_2-\text{COOH}$, p-aminophenyl alanine, $\text{H}_2\text{N}-\text{F}(\text{NH})-\text{NH}-(\text{C}_6\text{H}_6)-\text{CH}_2-\text{COOH}$, p-guanidinophenyl alanine or pyridinoalanine (Pal). These latter residues may form hydrogen bonding with the OH^- moieties of the Tyrosine residues at the MHC-1 N-terminal binding pocket, as well as to create, at the same time aromatic-aromatic interactions.

Side chains of amino acid derivatives at positions P4-P8 (if these are Ser, Tyr, Lys, Cys or Orn can be protected-attached to alkyl, aryl, alkylol or arylol moieties. These groups may enhance the binding to the T cell receptor.

Longer derivatives in which the second anchor amino acid is at a P10 position may include at P9 most L and D amino acids. Accommodation of longer peptides is possible through zig-zag or bulge modes. In some cases shorter derivatives are also applicable.

Cyclic derivatives should include preferably amino acids at position P4, P5, P6, and P7. Cyclization can be obtained through amide bond formation, e.g., by incorporating Glu, Asp, Lys, Orn, di-amino butyric (Dab) acid, di-

aminopropionic (Dap) acid at various positions in the chain (-CO-NH or -NH-CO bonds). Backbone to backbone cyclization can also be obtained through incorporation of modified amino acids of the formulas $\text{H-N}((\text{CH}_2)_n\text{-COOH})\text{-C(R)H-COOH}$ or $\text{H-N}((\text{CH}_2)_n\text{-COOH})\text{-C(R)H-NH}_2$, wherein $n = 1-4$, and further wherein R is any natural or non-natural side chain of an amino acid.

Cyclization via formation of S-S bonds through incorporation of two Cys residues is also possible. Additional side-chain to side chain cyclization can be obtained via formation of an interaction bond of the formula $\text{-(CH}_2\text{)}_n\text{-S-CH}_2\text{-C-}$, wherein $n = 1$ or 2 , which is possible, for example, through incorporation of Cys or homoCys and reaction of its free SH group with, e.g., bromoacetylated Lys, Orn, Dab or Dap.

Peptide bonds (-CO-NH-) within the peptide may be substituted by N-methylated bonds (-N(CH₃)-CO-), ester bonds (-C(R)H-C-O-O-C(R)-N-), ketomethylen bonds (-CO-CH₂-), α -aza bonds (-NH-N(R)-CO-), wherein R is any alkyl, e.g., methyl, carba bonds (-CH₂-NH-), hydroxyethylene bonds (-CH(OH)-CH₂-), thioamide bonds (-CS-NH-), olefinic double bonds (-CH=CH-), retro amide bonds (-NH-CO-), peptide derivatives (-N(R)-CH₂-CO-), wherein R is the "normal" side chain, naturally presented on the carbon atom.

These modifications can occur at any of the bonds along the peptide chain and even at several (2-3) at the same time. Preferably, but not in all cases necessary, these modifications should exclude anchor amino acids.

Natural aromatic amino acids, Trp, Tyr and Phe, may be substituted for synthetic non-natural acid such as TIC, naphthylelanine (Nol), ring-methylated derivatives of Phe, halogenated derivatives of Phe or o-methyl-Tyr.

As used herein in the specification and in the claims section below, the term "transduced" refers to the result of a process of inserting nucleic acids into cells. The insertion may, for example, be effected by transformation, viral infection, injection, transfection, gene bombardment, electroporation or any other means effective in introducing nucleic acids into cells. Following transduction the nucleic acid is either integrated in all or part, to the cell's genome (DNA), or remains external to the cell's genome, thereby providing stably transduced or transiently transduced cells.

As used herein in the specification and in the claims section below the phrase "derived from a protein" refers to peptides derived from the specified protein or proteins and further to homologous peptides derived from equivalent regions of proteins homologous to the specified proteins of the same or other species, provided that these peptides are effective as anti-tumor vaccines. The term further relates to permissible amino acid alterations and peptidomimetics

designed based on the amino acid sequence of the specified proteins or their homologous proteins.

As used herein in the specification and in the claims section below the phrase "anti-tumor vaccines" refers to a vaccines effective in preventing the development of, or curing, cancer, including primary tumor and/or metastases.

As used herein in the specification and in the claims section below the phrase "prevention or cure" also refers to inhibiting, slowing or reversing the progression of a disease, substantially ameliorating clinical symptoms of a disease or substantially preventing the appearance of clinical symptoms of a disease.

According to a preferred embodiment of the present invention the Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1) proteins are each independently of a mammal, e.g., a humanoid such as a human being or a rodent such as murine.

According to another preferred embodiment of the present invention each of the one or more tumor associated antigen peptides includes a number of residues, e.g., 7, 8, 9, 10 and 11 residues, including 2-3 residues which form a primary anchor for interaction with MHC class I molecules and 2-3 residues which bind to T cell receptors.

According to yet another preferred embodiment of the present invention the composition further comprising a carrier, wherein the tumor associated antigen peptide(s) are presented in context of the carrier.

The carrier can be a proteinaceous carrier to which the peptides are linked. Methods of linking short peptides to carriers are well known in the art of vaccination. The carrier can alternatively be a particulate adjuvant, an oil or emulsifier based adjuvant, a gel based type adjuvant, or an adjuvant based on specific targeting of antigen, such as, but not limited to, antibody-liposome conjugates. The carrier can also be a protein or a recombinant protein produced, for example in bacteria, yeast or in mammalian cells, including cytokines, such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, interferon alpha, interferon beta, interferon gamma and others. The carrier can also be an antigen presenting cell, such as, but not limited to, a dendritic cell, a macrophage, a B cell or a fibroblast. The cell is selected either autologous or non-autologous HLA matching cell. Optionally the cell can be a cultured cell, a treated cell by various reagents (e.g., by early and/or late acting cytokines), transduced by genes, and/or irradiated or radiated.

The vaccine composition according to the present invention is effective in prevention or cure of cancer and/or cancer metastases. In other words, the composition is effective for primary tumors, secondary tumors and metastases thereof in the same organ or in another organ, provided that the tumor expresses the above listed tumor associated proteins. According to a preferred embodiment of the present invention the cancer being treated or prevented via the administration of the vaccine composition is a carcinoma. i.e., a malignant tumor composed of epithelial tissue. The cancer treated or prevented according to the present invention can be, for example, breast, bladder, prostate, pancreas, ovary, thyroid, melanoma, colon, stomach and/or head and neck cancer.

According to an embodiment of the present invention, the Uroplakin protein can be Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.

According to another embodiment of the present invention, the tumor associated antigen peptides derived from the Mucin protein are from a non tandem repeat array of Mucin.

According to yet another embodiment of the present invention, one or more tumor associated antigen peptides derived from the Mucin protein are from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of Mucin.

According to a presently preferred embodiment of the present invention the one or more tumor associated antigen peptides are selected from SEQ ID NOs: 1-49 and effective homologues and analogs thereof.

For therapeutic or prophylactic anti-tumor treatment, the vaccine composition according to the present invention may include thickeners, carriers, buffers, diluents, surface active agents, preservatives, and the like, all as well known in the art. The composition may also include one or more active ingredients, such as, but not limited to, anti-inflammatory agents, anti-microbial agents, anesthetics and the like.

The vaccine composition may be administered in either one or more of ways. Administration may be effected topically (including ophtalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip or intraperitoneal, subcutaneous, or intramuscular injection.

Compositions for topical administration can include, but are not limited to, lotions, ointments, gels, creams, suppositories, drops, liquids, sprays and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

Formulations for parenteral administration can include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents, adjuvant and other suitable additives. The adjuvant is preferably of a type allowed for use in treating human beings, such as BCG adjuvant.

5 Dosing is dependent on responsiveness, but will normally be one or more doses per week or month, with course of treatment lasting from several weeks to several months. Persons ordinarily skilled in the art can easily determine optimum dosages, dosing methodologies and repetition rates.

10 The present invention provides novel tumor associated antigen peptides effective in eliciting CTL response which can therefore be effective therapeutic agents to combat cancer.

Each of the various and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the Examples section that follows.

15

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

20

MATERIALS AND EXPERIMENTAL METHODS

Mice: The derivation of HLA-A2.1/D^b- β 2m monochain, transgenics, H-2D^b x β 2m double knockout mice (HhD mice) was described before (55).

25 **Tumor and cell lines:** RMA-S is a TAP-2 deficient lymphoma clone of C57BL/6 origin. The RMA-S HhD and RMA-S-HhD-B7.1 are transfectants carrying the HhD construct without or with the gene for murine B7-1 costimulatory molecule, respectively. T2 is a TAP-2 deficient human lymphoblastoid line expressing HLA-A2. RMA-S, RMA-S derivatives and T2
30 cells were maintained in RPMI 1640-10 % FCS and antibiotics.

The human tumor cell lines, J82 (TCC of the bladder) sup (TCC of the bladder, DU145 (prostate carcinoma) and MDA-MB-157 (breast carcinoma) and the HhD transfectants of DU145 and MDA-MB-157 were maintained in DMEM-10 % FCS 2 mM glutamine - 1 mM sodium pyruvate - 1 % non-essential amino
35 acids and antibiotics.

HhD and B7-1 transfectants were grown in presence of 500-1000 μ g/ml of Genitcin, G418. All tissue culture media and supplements were purchased from Gibco.

Peptide synthesis: Peptides were synthesized on an ABIMED AMS 422 multiple peptide synthesizer (Langenfeld, Germany), employing the α -N-9-fluorenylmethoxycarbonyl (Fmoc) strategy following the company's commercially available protocols. Peptide chain assembly was conducted on a 2-chlorotrityl chloride resin (Novabiochem, Laufelfingen, Switzerland). Crude peptides were purified to homogeneity by reversed-phase HPLC on a semi-preparative silica C-8 column (250 x 10 mm; Lichnonorb RP-8; Merck). Elution was accomplished by a linear gradient established between 0.1 % trifluoroacetic acid in water and 0.1 % trifluoroacetic acid in 70 % acetonitrile in water (v/v). Purity of peptides (\cong 96 %) was ascertained by analytical HPLC (RP-18) using various acetonitrile water (containing 0.1 % trifluoroacetic acid) gradients. The products' compositions were determined by amino acid analysis (Dionex automatic amino acid analyzer, Sunnyvale, CA) after extraction acid hydrolysis. Molecular weight was ascertained by mass spectrometry (VG Tofspec; Laser Desorption Mass. Spectrometry; Fison Instruments, Manchester, U.K.). Tables 1-8 present these peptides.

Preparation of peptide fractions from fresh human tumors and normal tissues: Surgical specimens of TCC, normal bladder mucosa, prostate carcinoma, hyperplastic benign prostate, breast carcinoma or normal breast tissue were transferred on ice from surgery to a qualified pathologist. Tumor tissue were cleaned from most adjacent tissue and snap frozen. Non-necrotic fragments, from 5-10 patients were homogenized in PBS; 0.5 % Nonidet P-40; 10 μ g/ml soybean trypsin inhibitor; 5 μ g/ml lupeptide; 8 μ g/ml aprotonine and 0.5 mM PMSF and homogenized using a glass-teflon homogenizer. Following further stirring for 30 minutes at 4 °C, the homogenates were titrated with 10 % TFA to a final concentration of 0.1 % TFA and stirred for 30 minutes at 4 °C. After ultra-centrifugation for 30 minutes at 42 Krpm the supernatants were applied to Sephadex G25 columns and fractions were monitored at 230 nm. Peptide fractions under 10 Kd were pooled, lyophilized and further fractionated by centriprep 3 by centrifugation (Amicon, Beverly, MA). The peptide pool after lyophilization, was dissolved in opti-MEM (Gibco) for further use.

Scoring of HLA-A2.1 binding peptides: Protein sequences were screened for MHC binding by a worldwide web interface at http://www.bimas.dcrt.nih.gov/molbio/hla_bind/index.html. The software, based on accumulated data scores every possible peptide in the protein for possible binding to MHC according to the contribution of every amino acid in the peptide. Theoretical binding scores represent calculated half-life of the HLA-A2.1-peptide complex.

Measurement of peptide binding by stabilization of cell surface MHC:

Peptide binding to HhD was measured by stabilization of HhD on RMA-S transfectants, using an indirect FACS assay as follows: 5×10^5 peptide loaded TAP-2 deficient RMA-S-HhD cells (see vaccination), were incubated with anti
 5 HLA monoclonal antibodies for 30 minutes at 4 °C. After washing the cells with PBS - 0.5 %, BSA + 0.1 %, sodium azide, the second antibody, goat anti mouse-FITC (Jackson Lab.), was applied for 30 minutes at 4 °C. Following washing, the amount of bound antibodies was detected by FACSScan (Bacson).

Mouse monoclonal antibodies B-9-12, w6/32 (anti HLA A, B, C) and
 10 B.B7.2 (anti HLA-A2.1) were used for analyses.

Vaccination: Mice were immunized intraperitoneally three times at 7-day intervals, with 2×10^6 irradiated (5,000 rad) tumor cells, or with peptide-loaded RMA-S-HhD-B7.1 transfectants. Peptide loading of RMA-S-HhD-B7.1 cells was performed as follows: After washing the cells 3 times in PBS, the surface
 15 expression of HhD monochins was stabilized by a 4 hours culture at 26 °C. Synthetic peptides, or peptide extracts were added to 10×10^6 cells in 1 ml of opti-MEM (Gibco) medium to a concentration of 100 μ M or 1 mM, respectively. The cells were incubated at 26 °C and for additional 3 hours at 37 °C. Peptide loaded RMA-S-HhD - B7.1 cells were irradiated (5000 rad), washed,
 20 resuspended in PBS and injected into mice. In mixed synthetic vaccines, RMA-S-HhD-B7.1 cells were loaded separately with each peptide and pooled before vaccination.

In vitro cytotoxicity assays: Mice were immunized intraperitoneally three times at 7-day intervals, with 2×10^6 irradiated (5,000 rad) tumor cells, or with
 25 peptide-loaded RMA-S-HhD-B7.1 transfectants. Spleens were removed on day 10 after the last immunization and splenocytes were restimulated *in vitro* with either irradiated tumor cells (for mice immunized with tumor cells) or with lymphocytes pulsed with 100 μ M synthetic peptides or 1 mM patients-derived extract in opti-MEM (Gibco) medium. Restimulated lymphocytes were
 30 maintained in RPMI 1640 medium containing 10 % FCS, 1 mM glutamine, combined antibiotics, 1 mM sodium pyruvate, 10 mM HEPES, pH 7.4, 5×10^{-5} M β -mercaptoethanol, and 1 % nonessential amino acids (RPMI-HEPES medium) for 5 days. Viable lymphocytes (effector cells), were separated by lymphocyte-M (Cedarlane, Ontario, Canada) centrifugation, resuspended in
 35 RPMI-HEPES medium, and admixed at different ratios with 5×10^3 peptide loaded RMA-S-HhD cells, previously loaded with different peptides (see vaccination). CTL assays were performed in U-shaped microtiter wells, at 37 °C, 5 % CO₂ for 5 hours. Cultures were terminated by centrifugation at 1,000 rpm

for 10 minutes at 4 °C. A total of 100 µl of the supernatants was mixed with scintillation fluids and counted in a beta counter. Percentage of specific lysis was calculated as follows: % lysis = (cpm in experimental well - cpm spontaneous release)/(cpm maximal release - cpm spontaneous release) x 100. Spontaneous release was determined by incubation of 100 µl-labeled target cells with 100 µl of medium. Maximal release was determined by solubilization of target cells in 100 µl 0.1 M NaOH.

Sensitization of human PBL *in vitro*: Patients were HLA typed by the Tissue Typing Unit of The Rabin Medical Center, Israel. Only HLA-A2.1 carrying haplotypes were used herein. Heparanized blood, 40 ml/patient was separated on Ficoll gradients, mononuclear cells were collected at the interphase, washed three times in PBS and resuspended in PBS. A third of the lymphocyte volume was incubated with tumor extracted peptides, at a concentration of 1 O.D230/ml for 2 hours at 37 °C. The other two thirds of the cells were suspended at 10⁶ cells/ml in RPMI - 10 % FCS, 2 mM glutamine, 1 mM sodium pyruvate, 10 mM Hepes, 5 x 10⁻³ M β-mercaptoethanol and combined antibiotics. Peptide loaded cells were added together with 5-20 u/ml of recombinant human IL-2 (Peptotec, Rehovot, Israel). Cells were incubated for 5-7 days at 37 °C, 5 % CO₂ and then used in CTL assays as described above.

EXPERIMENTAL RESULTS

EXAMPLE 1

Uroplakins are T cell defined TAAs in transitional cell carcinoma (TCC)

Transitional cell carcinoma (TCC) of the bladder represents a prevalent cancer. Its incidence increases gradually during life and peaks in the 7th and 8th decade to 200/100,000. About 95 % of bladder tumors are TCC. Of these, 75 % are superficial papillary tumors and have a 15 % - 20 % progression rate during the first two years of follow up. Approximately, 20 % - 25 % of TCC present as muscle invasive tumors with aggressive behavior and a 40 % survival rate at 5 years (56). After surgical resection of superficial tumors, intravesical instillations of chemotherapeutics as Tiotepa or Mitomycin, or non-specific intravesical immunotherapy with Bacillus Calmette Guerein (BCG), reduce the frequency of recurrences and have a minimal effect on progression. BCG intravesical instillation was found to be the most effective in reducing the recurrence and progression rate of high grade papillary TCC and of transitional cell carcinoma *in situ*. The mechanism of action was not yet elucidated. However, following

therapy there is a mononuclear infiltrate of the lamina propria of the bladder mucosa. This infiltrate is rich in T-cells, macrophages and Langerhans cells (57). It was found that CD8 T-cells were activated after BCG intravesical therapy and induced specific lysis of TCC cells (58). Also, BCG intravesical instillation
 5 resulted in increased presentation of MHC complexes on transitional cell membrane (59). Lately, CTL, which induced specific lysis of autologous tumor cells, were identified among tumor infiltrating lymphocytes grown from transitional cell carcinomas of the bladder (60). It seems that specific cytotoxic T-cell immunity against TCC may be augmented by BCG intravesical therapy.

10 An important group of TAAs are actually differentiation antigens. Such antigens were shown to induce CTL in melanoma and in colon carcinoma patients (37, 38).

Transitional epithelium is a differentiated multilayered epithelium which is restricted to the urinary tract. It is composed of 4 to 7 cell layers: a basal cell
 15 layer; an intermediate 2-3 cell layer and an apical (luminal) layer composed of umbrella like cells. These latter cells are connected by tight junctions and create an impermeable barrier to water and urinary solutes.

Lately, differentiation proteins of transitional epithelium have been characterized morphologically and biochemically. These proteins were found in
 20 urothelial plaques which are present in the luminal plasma membrane of urothelial superficial (umbrella) cells. The molecular constituents of these plaques comprise four transmembranal proteins designated Uroplakins (UP); UP Ia (47 Kd); UP Ib (28 Kd); UP II (15 Kd) and UP III (47 Kd glycoprotein; the size of the core protein is 28.9 Kd). Uroplakin III probably plays a role in the
 25 formation of the urothelial glycocalyx and may interact with the cytoskeleton. An extensive evaluation of UP tissue distribution showed that UP are restricted to normal transitional cell epithelium and to TCC (61). In normal transitional epithelium the UPs were found in the apical membrane of the superficial umbrella cells. In superficial papillary TCC, the UPs were localized in the apical
 30 membrane of the luminal cells and in the cell membranes lining intra and inter-cellular lumina. In invasive TCC, UPs were localized more randomly in the cell membrane of the cells which infiltrated the stroma. Recently, the sequence of human Uroplakin II cDNA and Uroplakin Ia genomic sequence were determined and their protein sequences were deduced (NCBI accession Nos. 2190407 and
 35 2098577, respectively). Since TCC patients treated by BCG were shown to have increased T cell infiltrates in the bladder mucosa, we first tested whether BCG treated patients show increased CTL activity against TCC derived peptides and whether UP II peptides may constitute CTL specific epitopes. We further used

the TCC model to test the T cell repertoire overlap between patient derived CTL and CTL induced in the unique H-2D^b/- x β 2m/- x HhD^{+/+} (HhD) mice.

Cytotoxic activity of human peripheral blood lymphocytes (PBL) against target cells pulsed with Uroplakin II peptide homologues: The human UP II amino acid sequence was screened for potential HLA-A2 binding, 9 amino acids peptides by a HLA binding motif program. We chose that particular HLA allele because it is present in 45 % of the population. Seven 9-mers peptides, which were predicted to bind with high affinity to HLA-A2 were selected and synthesized (Table 1). The binding of these peptides to HLA-A2 was tested by an MHC stabilization assay utilizing TAP negative cells, which can present exogenously bound peptides. The RMA-S cells (55) transfected by the HhD construct were loaded with various concentrations of UP II peptides and then reacted with anti-HLA antibody. High affinity binding was shown for all 7 peptides by FACS analyses (Figure 1).

TABLE 1

Predicted human Uroplakin II peptides that bind to HLA-A2

Peptide	Sequence	Amino-acids*	SEQ ID:
HUO1	FLLVLGFII	168-176	1
HUO2	VLPSVAMFL	161-169	2
HUO3	LVLGFIIAL	170-178	3
HUO4	KVVTSSFVV	67-77	4
HUO5	LVPGTKFYI	109-117	5
HUO6	LLPIRTLPL	4-12	6
HUO7	YLVKKGTAT	119-127	7

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the peptide in the protein are listed. * Numbering is according to NCBI accession No. 2190407.

To restimulate CTL activity in patient's PBL we chose 6 TCC patients of the HLA-A2 haplotypes: 4 patients underwent adjuvant intravesical BCG instillations and 2 underwent no adjuvant therapy after endoscopic resection of the TCC. Another patient without TCC served as control. PBL were stimulated *in vitro* by a mixture of tumor acid extracted peptides from 9 TCC samples as described before (62). Each tumor was extracted individually and an equimolar ratio of each sample was added to the mixture. The final concentration of

peptides in the stimulation reaction was 1 O.D. 230/ml. Antigen presenting cells (APC) from the same peripheral blood were pulsed with the tumor extract. Peripheral blood lymphocytes (PBL) were stimulated with APC loaded with pooled TCC extract and with IL2 for 5 days. Uroplakin II peptides, TCC
 5 extracted peptides and normal bladder extracted peptides were loaded on labeled T2 cells (HLA-A2 positive cells which are TAP deficient, express empty MHC molecules and can be loaded only with exogenous peptides), and were screened by CTL obtained from PBL of the patients, as described. CTL from TCC patients treated with BCG induced 32 % to 71 % specific lysis of Uroplakin II
 10 peptide loaded T2 cells (Figure 2). Tumor extract loaded targets were lysed at 62 % while normal bladder extract loaded targets were lysed at 29 % only, indicating preferential recognition of tumor antigens. CTL from TCC patients without BCG adjuvant therapy induced 25 % to 48 % specific lysis of T2 cells loaded Uroplakin II peptides, 29 % lysis of tumor extract loaded T2 cells and 15
 15 % lysis of bladder extract loaded T2 cells. There was no lysis of T2 cells loaded with breast tumor extract, or with a HLA-A2 binding peptide homologue to tyrosinase, a melanoma differentiation antigen. A TCC cell line expressing HLA-A2, J82 was effectively killed while another TCC line expressing low levels of HLA, SUP, was not lysed. Also, there was no lysis of a natural killer
 20 (NK) sensitive cell line (K562). CTL obtained from the patient without TCC induced no specific lysis of tumor extract, or Uroplakin II peptides loaded APC cells. The differences between TCC patients treated with BCG, TCC patients without adjuvant therapy and non-TCC patients were very significant ($P < 0.001$). We also tested the activity of patient's PBL, resensitized by the pool of
 25 tumor extracts, against individual tumor extracts. Figure 3 shows higher activity of lymphocytes derived from BCG treated patients than lymphocytes derived from non-treated patients. Control lymphocytes show low activity. Thus shared antigens are expressed in all TCC samples.

CTL induced in HhD mice recognize the same UP II peptides as
 30 **human PBL:** The immunogenicity of Uroplakin homologue peptides was also evaluated in a mouse model, in which the murine MHC class I genes are not expressed ($Db^{-/-}$, $\beta 2$ -microglobulin $-/-$, double knockout) and which is transgenic for a HLA-A2 and human $\beta 2$ -microglobulin monochain (HhD mice). The advantage of a HLA-A2 transgenic mouse model is that CTL epitopes can be
 35 detected more easily and reproducibly, without lengthy and repeated *in vitro* stimulations. The murine MHC knockout mice used herein has the additional advantage that their CTL repertoire is only HLA-A2 restricted and the CTL response against human tumor extract or human cell lines is not masked by a

potential xenogeneic response. HhD mice were immunized with the 7 Uroplakin II homologue peptides or with the same TCC peptide extract used for stimulation of patient's PBL. Lymphocytes were obtained from the spleens of these mice and restimulated *in vitro* once with the corresponding peptides.

5 The 7 UP II peptides, that were recognized by human CTL, 8 new UP Ia homologues peptides with high HLA-A2 binding affinity (Table 2), 4 murine UP II derived HLA-A2 binding peptides (Table 3), TCC and normal bladder mucosa peptide extracts and two TCC cell lines: J82 (HLA-A2 positive) and TCCSUP (minimal expression of HLA-A2) were screened by the CTL obtained from these
10 mice.

TABLE 2

Predicted human Uroplakin Ia peptides that bind to HLA-A2

	Peptide	Sequence	Amino-acids*	SEQ ID:
15	HUO11	SLFAETIWV	33-41	8
	HUO12	MLIAMYFYT	248-256	9
	HUO13	LMWTLPVML	241-249	10
	HUO14	MLIVYIFEC	100-108	11
	HUO15	YIFECASCI	104-112	12
20	HUO16	LVLMLIVYI	97-105	13
	HUO17	ALCRRRSMV	85-93	14
	HUO18	LLSGLSLFA	28-36	15

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter
25 amino acid codes and the position of the peptide in the protein are listed. * Numbering is according to NCBI accession No. 2098577.

TABLE 3

Predicted murine Uroplakin II peptides that bind to HLA-A2

	Peptide	Sequence	Amino-acids*	SEQ ID:
30	MURO1	FLLVVGLIV	168-176	16
	MURO3	LVVGLIVAL	170-178	17
	MURO4	KVVKSDFFV	69-77	18
	MURO6	TLPVQTLPL	4-12	19

35 The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the peptide in the protein are listed. The murine 1, 3, 4, 6 peptides are homologous to the human Uroplakin II peptides 1, 3, 4, 6 in Table 1. * Numbering is according to NCBI accession No. 586160.

The results are expressed as lytic units (LU30), which represent 10^6 /the number of effectors which induce 30 % specific target lysis. More LU signify a more potent CTL response. Figure 4 shows that CTL from HhD mice immunized with Uroplakin II peptides (1-7 loaded individually and injected by a mixture of loaded cells) induced a significant lysis of RMA-S-HhD cells loaded with Uroplakin II homologue peptides (8.1 to 32.2 LU). There was no lysis of RMA-S-HhD cells loaded with nonspecific peptides as breast tumor extract or the HLA-A2 binding tyrosinase homologue peptide. Also, there was no lysis of RMA-S-HhD cells without peptides (empty target cells). There was intense lysis of J82 cells (17.3 LU) and there was much less lysis of TCCSUP cells (1.2 LU). There was also intense lysis of RMA-S-HhD cells loaded with TCC peptide extract (15 LU) and less lysis of normal bladder mucosa peptide extract loaded cells (4.5 LU) (Figure 4).

CTL from mice immunized with TCC peptide extract induced significant lysis of RMA-S-HhD targets loaded with Uroplakin II homologue peptides (3.2 to 14 LU); Uroplakin Ia homologue peptides (2 to 11.4 LU); TCC peptide extract (12.6 LU) and less lysis of these targets loaded with normal bladder mucosa peptide extract (2 LU). There was also intense lysis of J82 cells (7.6 LU) and less lysis of TCCSUP cells (0.8 LU). There was no lysis of target cells loaded with nonspecific peptides or of empty target cells. The lysis pattern of the targets loaded with the different Uroplakin II homologue peptides was similar to the CTL assays using human PBL, CTL from HhD mice immunized with Uroplakin II peptides or with TCC peptide extract (Figure 5).

Although human and HhD murine CTL recognize the same UP II peptides, it was not clear whether HhD murine CTL are directed against the actual T cell epitopes or against possible differences between humanoid and rodent UP II. We evaluated also the crossreactivity of CTL from mice immunized with human Uroplakin II peptides against 4 homologue murine Uroplakin II peptides. The murine UP II sequence was screened for HLA-A2 binding peptides and 4 peptide homologues to human peptides 1, 3, 4 and 6 were synthesized. Murine peptides differ in 1-3 amino acids from similar human peptides, yet most are conservative changes (Tables 1 and 3). Murine UP II peptides bind stably to RMA-S-HhD cells.

There was significant lysis of the targets loaded with each of these murine peptides. A good correlation between the LU30 results for the human and their homologue murine peptides was found. Three human peptides induced more target lysis than their murine counterpart and for the fourth peptide the trend was opposite (Figures 4 and 5). Thus, the specificity of the HhD murine CTL, as the

specificity of human PBL is directed to common epitopes on the peptides. To examine whether any autoimmune effects are observed in peptide vaccinated HhD mice, the internal organs of the HhD mice immunized with TCC peptide extract or with Uroplakin II homologue peptides were formalin fixed and representative sections from each organ were stained with hematoxylin and eosin. Two non-immunized HhD mice served as control. There were no pathological finding such as tissue necrosis or inflammatory infiltrate in any of the organ examined. The urinary bladder was carefully examined using a larger number of sections. The bladder wall including the mucosa was normal in the immunized and in the control mice.

EXAMPLE 2

PSA, PSMA and PAP homologue peptides are immunogenic CTL epitopes in HLA-A2 transgenic H-2D^b, β 2m double knockout mice

Carcinoma of the prostate (CAP) is the most prevalent cancer and the second cause of death in man (63). The use of prostate specific antigen (PSA) in early detection of CAP has caused a stage shift of the disease (64). By now, 70 % of CAP cases are detected when they are still organ confined. These patients may be cured by radical prostatectomy or by radiation therapy. However, the long term cancer specific survival of organ confined disease after radical prostatectomy, even in the best series, is only 70 % (65). Patients with large tumors, high grade tumors and those with seminal vesicle invasion, or positive lymph nodes are at increased risk of metastases and death due to CAP (66). However, CAP is a slow growing tumor and there is no effective adjuvant chemotherapy (67). Androgen ablation by surgical or medical means does not increase survival, but only increase time to recurrence and palliate symptomatic bone metastases (68). There is a need for an effective adjuvant or neo-adjuvant therapy for CAP and for systemic therapy for metastatic CAP. Thus, specific immunity is being investigated as a potential adjuvant therapy. PSA and prostate specific membrane antigen (PSMA) are proteins which are expressed almost exclusively in prostate tissue, benign or malignant and are differentiation antigens. Prostate acid phosphatase (PAP) is expressed also in other tissues but its concentration is much higher in prostate tissue. In a few studies, 3 PSA homologue peptides (69-70) and 2 PSMA homologue peptides (71-72) were found to be immunogenic and to induce specific CTL by repeated *in vitro* stimulation of peripheral blood lymphocytes from CAP patients. In one of these studies (72), patients with metastatic prostate cancer were immunized with the 2

immunogenic PSMA peptides, loaded on dendritic cells. There was some partial responses expressed by a reduction in PSA levels and some objective reduction in bone metastases burden. However, the procedures used to induce CTL lines *in vitro* are cumbersome, lengthy and not reliable. Some CTL peptide epitopes which bind with high affinity to MHC class I may induce tolerance and will be not detected. Moreover, repeated *in vitro* stimulations, necessary for inducing CTL lines may select CTL which have a growth advantage in culture and not the most potent and specific CTL *in vivo*. Herein the HLA-A2 transgenic, H-2D^b β 2 microglobulin double knockout mice (HhD mice) was used for identification of novel prostate cancer specific CTL epitopes. Peptides homologue to PSA, PSMA and PAP, which are expressed mainly in benign or malignant prostate tissue, were evaluated for their capacity to induce specific CTL in HhD mice. The amino acid sequences of these proteins (NCBI accession Nos. 296671, 2897946 and 439658) were screened by a HLA binding motif program. The peptides which were predicted to bind to HLA-A2 were synthesized (Tables 4, 5, 6) and their binding affinity was tested on RMA-S-HhD cells (TAP deficient cells, capable of presenting only exogenous peptides and which were transfected with the HhD construct) by FACS (Figures 6 and 7). The peptides with high binding affinity were used for immunization of HhD mice. Five PSA homologue peptides, in addition to the peptides detected by other authors, 6 PSMA homologue peptides, including the 2 peptides detected by other authors (PSMA 5 and 6) and 4 PAP homologue peptides were found.

TABLE 4

Predicted human prostate specific antigen (PSA) peptides that bind to HLA-A2

Peptide	Sequence	Amino-acids*	SEQ ID:
PSA1	DLHVISNDV	175-183	20
PSA2	VLVHPQWVL	53-61	21
PSA3	FLRPGDDSS	110-118	22
PSA4	ALGTTCYAS	147-155	23
PSA5	KLQCVDLHV	170-178	24

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the peptide in the protein are listed. * Numbering is according to NCBI accession No. 296671.

TABLE 5

Predicted human prostate specific membrane antigen (PSMA) peptides that bind to HLA-A2

	Peptide	Sequence	Amino-acids*	SEQ ID:
5	PSMA1	ELAHYDVLL	109-117	25
	PSMA2	NLNGAGDPL	260-268	26
	PSMA3	TLRVDCITFL	461-469	27
	PSMA4	MMNDQLMFL	663-671	28
	PSMA5	ALFDIESKV	711-719	29
10	PSMA6	LLHETDSAV	4-12	30

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the peptide in the protein are listed. * Numbering is according to NCBI accession No. 2897946.

15

TABLE 6

Predicted human prostate acid phosphatase (PAP) peptides that bind to HLA-A2

	Peptide	Sequence	Amino-acids*	SEQ ID:
20	PAP1	VLAKELKFV	30-38	31
	PAP2	ILLWQPIPV	135-143	32
	PAP3	DLFGIWSKV	201-209	33
	PAP4	PLERFAELV	352-360	34

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the peptide in the protein are listed. * Numbering is according to NCBI accession No. 439658.

HhD mice were immunized and boosted twice with the PSA, PSMA, or PAP peptides. Peptides were loaded individually on RMA-S-HhD cells and equal numbers of pulsed cells were mixed in the vaccine. Mice were also immunized with CAP peptide extract, or with DU145-HhD, a CAP cell line transfected with the HhD construct. Lymphocytes were obtained from the spleen of these mice and restimulated once, *in vitro* with the corresponding peptide tumor extract or cells. PSA, PSMA, PAP, CAP and normal prostate peptide extract and DU145-HhD cells were screened by the CTL obtained from these mice. The results are expressed as lytic units (LU) which represent 10^6 /Number of CTL effectors which induce 30 % specific target lysis. More lytic units signify a more potent CTL response.

CTL from HhD mice immunized with PAP peptides induced intense lysis of RMA-S-HhD cells (targets) loaded with PAP homologue peptides (20 to 65 LU, Figure 8). Three out of 5 PSA homologue peptides induced significant and specific lysis of targets loaded with the corresponding peptide (5.2 to 11.4 LU, Figure 9). Four out of 6 PSMA homologue peptides induced significant lysis of the targets loaded with the corresponding peptide (5 to 17 LU, Figure 10). Two out of these 4 peptides were found to be immunogenic by other authors, by *in vitro* stimulation of human peripheral blood lymphocytes. CTL obtained from each group of immunized HhD mice induced significant lysis of targets loaded with CAP peptide extract (4.5 to 22 LU). There was significantly less lysis of normal prostate peptide extract loaded targets than of the CAP peptide extract loaded targets for each immunizing peptide (Figures 8, 9 and 10). Also, there was intense lysis of DU-145-HhD cells by CTL obtained from HhD mice immunized with these cells (Figure 11), or immunized by PAP homologue peptides, or CAP peptide extract (28, 19 and 11 LU, respectively, Figure 12). There was no significant lysis of DU145-HhD cells by CTL derived from HhD mice immunized with PSA, or PSMA homologue peptides. These findings result probably from low expression of PSA and PSMA by DU145 cells.

There was good cross reactivity between CTL derived from mice immunized with CAP peptide extract and PSA, PSMA and PAP homologue peptides, which were found to be immunogenic in HhD mice. The CTL derived from mice immunized with DU145-HhD cells cross reacted with the immunogenic PAP homologue peptides (11 to 26 LU) CAP peptide extract (12 LU), and less with normal prostate peptide extract (3 LU). Overall, there was no lysis of targets loaded with nonspecific peptides such as breast cancer peptide extract, or HLA-A2 binding tyrosinase homologue peptide. Also, there was no lysis of RMA-S-HhD cells without peptides and there was no lysis of non transfected DU 145 cells (HLA-A2 negative). The internal organs of the HhD mice immunized with PSA, PSMA, PAP homologue peptides, or with CAP peptide extract, or with DU145-HhD cells, were formalin fixed and representative slices from each organ were stained with hematoxyllin and eosin. Two non-immunized HhD mice served as control. There was no pathological finding such as tissue necrosis or inflammatory infiltrate in any of the organ examined.

EXAMPLE 3

Breast specific CTL induction by BA46 (Lactadherin) peptides and by MUC1 peptides in HhD mice

5 Breast cancer is the second leading cause of cancer death among women in the Western World and the leading cause of death among women at the age of 30 to 70. Breast cancer afflicts 200,000 women per annum in the USA today. The highest mortality is restricted to patients whose regional lymph nodes are involved. Early detection, followed by surgery provides good prognosis. In
 10 patients with occult lymph node metastasis, adjuvant chemotherapy or hormonal therapy for breast cancer have been proven to be effective, yet a large fraction of patients will succumb to metastasis. (73).

As pointed out above, TAA vaccines may constitute an additional treatment modality for residual disease. A number of tumor associated antigens
 15 have been described for breast carcinomas. The MUC-1 Mucin, a high molecular weight glycoprotein is highly expressed on breast carcinomas. Specific, MHC-unrestricted recognition of MUC-1 by human cytotoxic T cells were demonstrated (74). More recently peptides from MUC-1 were shown also to induce MHC class I and MHC class II restricted responses (75, 76). HER2/neu
 20 derived peptides were shown to be recognized by CTL from breast carcinoma patients (77-78).

BA46 is a 46 kDa transmembrane-associated glycoprotein of the human milk fat globule membrane (HMFG), that is overexpressed in human breast carcinomas (79). The protein contains cell adhesion sequences (RGD), supports
 25 RGD based adhesion and interacts with integrin (80). It also contains an EGF-like domain, and a phospholipid binding sequence C1/C2-like domain of coagulation factor V and VIII. BA46 is present in the circulation of breast cancer patients but not in healthy individuals (81). Moreover, anti BA-46 radio-conjugated monoclonal antibodies, have successfully targeted human breast
 30 tumors transplanted into mice (82-83). Herein we tested MUC1 and BA46 derived peptides as potential TAA peptides in the HhD mouse system. BA46 homologous peptide vaccines induce HLA-A2 restricted CTL which preferentially recognize breast tumor derived peptides.

BA46 is overexpressed in many breast carcinomas, yet, no cellular
 35 immunity to BA46 protein has been reported so far. We evaluated peptides complementary to the amino acid sequences of human BA46, The sequence was screened for HLA-A2 binding motifs by a HLA binding motif program. Seven 9-

mer peptides which were predicted to bind with high affinity to HLA-A2 were selected and synthesized (Table 7).

TABLE 7

Predicted human breast associated BA46 peptides that bind to HLA-A2

Peptide	Sequence	Amino-acids*	SEQ ID:
EA46-1	KQGNFNAWV	271-279	35
BA46-2	NLLRRMWVT	131-139	36
BA46-3	NLFETPILA	356-364	37
BA46-4	NLFETPVEA	194-202	38
BA46-5	GLQHWVPEL	97-105	39
BA46-6	VQFVASYKV	313-321	40
BA46-7	RLLAALCGA	5-13	41

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the first amino acid of the peptide in the protein sequence as well as the calculated binding score are listed. * Numbering is according to NCBI accession No. 1589428.

All peptides bound well to the HhD molecules expressed on RMA-S transfectant (Figure 13). HhD mice were, vaccinated three times at weekly interval using 2×10^6 RMA-S-HhD/B7.1 with either tumor extract prepared from 5 samples of Breast carcinoma by acid extraction and separation of molecules < 3 Kd, or with a pool of BA46 peptides. Ten days after the last immunization, HhD spleen-derived antigen presenting cells (APC) were pulsed for 3 hour at 37 °C with either breast tumor extract or a BA46 peptide pool, followed by incubation with the rest of the splenocytes for 4 more days. BA46 peptides, breast extracted peptides and normal breast extracted peptides were loaded on labeled RMA-S-HhD cells and were screened by cytotoxic T lymphocytes (CTL) obtained from the HhD mice as described before (62).

Anti BA46 peptide CTL activity showed variability in immunogenicity at a range between 20 % to 60 %. Lysis of 60 % was obtained against BA46-7, 40 % against BA46-6, BA46-9 and 20 % against the rest of the peptides (Figure 14). CTL from HhD mice immunized with breast tumor extract peptide induced 10 % to 30 % specific lysis of BA46 peptide loaded RMA-S-HhD cells, 31 % lysis of tumor extract loaded targets and only 12 % and 14 % lysis of normal breast extracted peptides and colon extracted peptides loaded RMA-S-HhD cells, respectively. No lysis of the melanoma associated synthetic tyrosinase peptide was obtained (Figure 15). The ability of BA46 peptides to induce a breast

associated CTL reaction that is HLA-A2.1 restricted was further examined. CTL against individual BA46 peptides showed 30-50 % higher activity against breast tumor extract versus normal breast loaded target cells, supporting the fact that a preferential activity is obtained against breast tumor TAAS. These effectors also
 5 lysed preferentially a breast carcinoma line-HhD transfectant relative to parental cells, stressing the HLA-A2 restriction of the reaction (Figures 16 and 17).

EXAMPLE 4

10 Characterization of novel breast carcinoma MUC1 derived peptides as CTL epitopes

The polymorphic epithelial Mucin, encoded by the MUC-1 gene is a high molecular weight transmembranal glycoprotein overexpressed in a broad range of tumors, such as breast, pancreas, ovary, thyroid and myeloma (84-87) cancer. It
 15 was found that the growth rate of primary breast tumors induced by the polyoma middle T antigen is significantly slower in MUC-1 null mice, suggesting that MUC-1 might play a role in the progression of mammary carcinoma (88). Furthermore, high level of MUC-1 expression by human breast cancer cells was found to be directly correlated with the presence of axially lymph node metastasis
 20 (89). A major feature of the MUC-1 molecule is the presence of an highly immunogenic extracellular tandem repeat array (TRA) heavily O-glycosidic-linked serine and threonine residues (90-91). Altered carbohydrate structure of MUC-1 in breast cancer cells, is probably responsible for the exposure of core epitopes within MUC-1, specifically recognized by monoclonal antibodies, as
 25 well as non MHC-restricted cytotoxic T lymphocytes (92-93). HLA-A11 (94) and more recently for HLA-A2.1 (95) restricted responses to the extracellular TRA have also been reported. However it was suggested that Mucin can actively suppress cell-mediated response against glycosylated TRA (96), or induced direct T cell apoptosis (97). More recently Agrawal et al showed that synthetic peptides
 30 derived from MUC-1 TRA cause suppression of human T-cell proliferative responses (98). This data points out the ambiguous role of MUC-1 TRA in T cell activation.

A serie of peptides derived from non TRA domains of the MUC-1 as potential CTL epitopes. The MUC-1 sequence was screened for potential HLA-
 35 A2.1 binding peptides (Table 8) and the eight peptides were synthesized.

TABLE 8

Predicted human breast associated MUC-1 peptides that bind to HLA-A2

Peptide	Sequence	Amino-acids*	SEQ ID:
MUC1-/C6	LLLLTVLTV	31-40	42
5 MUC1-/D6	LLLTVLTVV	32-41	43
MUC1-/F6	FLSFHISNL	323-331	44
MUC1-/A5	LLVLVCVLV	442-451	45
MUC1-/F4	ALLVLVCVL	441-450	46
MUC1-/B5	SLSYTNPAV	519-528	47
10 MUC1-/A7	NLTISDVSV	412-421	48
MUC1-/E6	ALASTAPPV	226-234	49

The prediction program is described in Materials and Methods. The peptides are listed from the highest binder (top) to the lowest binder (bottom). Single letter amino acid codes and the position of the first amino acid of the peptide in the protein sequence as well as the calculated binding score are listed. * Numbering is according to NCBI accession No. 182253.

These 9 residue peptides are derived from the signal peptide, cytoplasmic and extracellular domains of the MUC-1 protein. Among these peptides only the MUC-1/B5 peptide has an identical sequence to the murine homologue. HLA-A2.1 binding of MUC-1- derived peptides were evaluation by FACS analysis. The selected peptides were loaded on the murine TAP-deficient RMA-S-HhD transfectants and MHC stabilization was monitored (Figure 18). Although all peptides bound efficiently at the 1-100 μ M range, 3 peptides MUC-1/D6, MUC-1/E6 and MUC-1/A7 exhibited higher binding affinity. In addition, similar binding affinities of these peptides were obtained upon loading on human TAP deficient T2 cells, expressing endogenous HLA-A2.1 molecules (data not shown).

Synthetic peptides corresponding the MUC-1 TRA epitopes were shown to induce CTL reaction in patients (94) as well as in HLA-A2.1K^b transgenic mice (95). Hence we primarily examined the lysis pattern of each of the particular peptides listed in Table 8 following an immunization with a pool of the MUC-1-derived synthetic peptides (Figure 19). CTL results showed significant lysis of RMA-S-HhD target cells loaded either with the MUC-1/D6 peptide (38 %) or with the MUC-1/A7 peptide (30 %). Lower lysis rate of 15 % with the MUC-1/E6 and with the MUC-1/F6 peptides was demonstrated. The rest of the peptides showed only background lysis properties. Hence MUC-1/D6, MUC-1/A7 and MUC-1/E6 harbor immunogenic properties.

To determine the processing and presentation of MUC-1-derived peptides by breast carcinoma tumors, we selected the MDA-MB-157 cell line which is characterized by high MUC-1 expression and low class I expression (data not shown). Both the parental tumor cell line and its HhD transfectant (MDA-MB-157-HhD), were used as targets in CTL assays (Figure 20). Mice were immunized with either RMA-S-HhD-B7.1 cells loaded with MUC-1 selected peptides MUC-1/D6, MUC-1/A7 and MUC-1/E6, or with peptide extract derived from fresh patients' tumors. Preferential lysis of the MDA-MB-157-HhD cell line by anti MUC-1-derived peptides activated lymphocytes suggested both breast associated as well as MHC-restricted lysis. Moreover, this data strongly support the processing and presentation of non TRA associated MUC-1-derived peptides in MDA-MB-157-HhD breast carcinomas cells. Further analysis showed inhibition of lysis by anti HLA monoclonal antibody w6/32 (data not shown). In addition specific lysis of MDA-MB-157-HhD cells by CTL directed against fresh tumor extracted peptides emphasized an overlap in the peptide repertoire between the breast tumor cell line and fresh breast tumor extract.

To prove the existence of MUC-1/D6, MUC-1/A7 and MUC-1/E6 peptides in patients' derived breast tumor peptide extracts, a CTL experiment was performed by utilizing CTL against breast tumor peptide extracts as effectors against target cells presenting MUC-1 peptides or a control of normal breast tissue (Figure 21). All 3 MUC-1-derived peptides, but not the HLA-A2.1 melanoma associated peptide tyrosinase, could be recognized and lysed by anti tumor extract CTL. RMA-S-HhD cells loaded with normal breast extract gave in different experiment between 40-50 % of the lysis induced against tumor extract loaded targets. This result is expected since a part of the tumor extract peptide repertoire consists of normal peptides.

A crucial parameter for selection of TAA peptides based vaccines are their expression frequency by tumors in comparison with normal tissues. Since MUC-1 protein is known to be overexpressed in tumors, with no tumor specific mutations, it is of obvious interest to examine the abundance of MUC-1 peptides in patients' derived normal breast tissue extract in comparison to tumor extract. CTL generated against MUC-1 peptides MUC-1/D6, MUC-1/A7 and MUC-1/E6, showed a 1.8-9.0 fold higher reactivity to tumor extract versus normal tissue extract (Figure 22). The same difference in preferential tumor versus normal tissue recognition by CTL could be detected upon vaccination with breast tumor extract peptides, supporting the window of specificity between normal and tumor tissues. These results suggest that MUC-1/D6, MUC-1/A7 and MUC-1/E6 are potential tumor associated antigen peptides.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended
5 to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

REFERENCES CITED

1. Fidler, I.J. and Balch, C.M. (1987) The biology of cancer metastasis and implications for therapy. *Curr. Progr. Surg.* **24**, 137-209. Boring, C.C., Squires, T.S., Tong, T., Montgomery, S.,
2. Abeloff, M.D. (1996) Breast Cancer. *Current opinion in oncology* **8**, 447-448.
3. Andriole, G.L. (1997) Adjuvant therapy for prostate cancer patients at high risk of recurrence following radical prostatectomy. *Eur. Urol. Supp* **3**; 65-69.
4. Nseyo, U.O. and Lamm, D.L. (1997) Immunotherapy of bladder cancer. *Semin. Surg. Oncol.* **13**, 342-349.
5. Kedar, E. and Klein, E. (1995) Cancer Immunotherapy: Are the results discouraging? Can they be improved? *Adv. Immunol.* **58**, 245-322.
6. Urban, J.L. and Schreiber, H. (1992) Tumor antigens. *Annu. Rev. Immunol.* **10**, 617-644.
7. Van Pel, A. and Boon, T. (1982) Protection against a nonimmunogenic mouse leukemia by an immunogenic variant obtained by mutagenesis. *Proc. Natl. Acad. Sci. USA* **79**, 4718-4722.
8. Pardoll, D.M. (1993) New strategies for enhancing the immunogenicity of tumors. *Curr. Opin. Immunol.* **5**, 719-725.
9. Qin, Z. and Blankenstein, T. (1996) Influence of local cytokines on tumor metastasis: using cytokine gene transfected tumor cells as experimental models. In "Attempts to Understand Metastasis Formation III", *Therapeutic Approaches for Metastasis Treatment, Current Topics in Microbiology and Immunology* (U. Gunlhert, P.M. Schlag and W. Birchmeier, eds.) **213III**, 55-64.
10. Eisenbach, L. (1996) Curing metastases? Gene and Peptide Therapy. *Ibid* pp. 85-100.
11. Gilboa, E. (1996). Immunotherapy of cancer with genetically modified tumor vaccines. *Semin oncol.* **23**, 101-107.
12. Williams, N. (1996) An immune boost to the war on cancer. *Science* **272**, 28-30.
13. Pardoll, D.M. (1996) Cancer vaccines: a road map for the next decade. *Curr. Opin. Immunol.* **8**, 619-621.
14. Boon, T. and van der Bruggen, P. (1996) Human tumor antigens recognized by T lymphocytes. *J. Exptl. Med.* **183**, 725-729.

15. Townsend, A. and Bodmer, H. (1989) Antigen recognition by class I-restricted T lymphocytes. *Annu. Rev. Immunol.* **7**, 601-624.
16. Rammensee, H.-G., Falk, K. and Rotzschke, O. (1993) Peptides naturally presented by MHC class I molecules. *Annu. Rev. Immunol.* **11**, 213-244.
17. Engelhard, V.H. (1994) Structure of peptides associated with class I and class II MHC molecules. *Annu. Rev. Immunol.* **12**, 181-207.
18. Chien, Y.H. and Davis, M.M. (1993) How alpha beta T-cell receptors 'see' peptide/ MHC complexes. *Immunol. Today* **14**, 597-602.
19. Feltkamp, M., Smits, H.L., Vierboom, M., Minnaar, R.P., deJough, B.M., Drijfhout, J.W., Schegget, J., Melief, C.J.M. and Kast, W.M. (1993) Vaccination with cytotoxic T lymphocyte epitope-containing peptide protects against a tumor induced by human papillomavirus type 16-transformed cells. *Eur. J. Immunol.* **23**, 2242-2249.
20. Rothbard, J.B. and Gefters, M.L. (1990) Interactions between immunogenic peptides and MHC proteins. *Annu. Rev. Immunol.* **9**, 527-566.
21. Treversari, C., van der Bruggen, P., Luescher, I.F., Lurquin, C., Chomez, P., Van Pel, A., de Plaen, E., Amar-Costesec, A. and Boon, T. (1992) A nonapeptide encoded by human MAGE-1 is recognized on HLA-A1 by cytotoxic T lymphocytes directed against tumor antigen MZ2-E. *J. Exp. Med.* **176**, 1453-1457.
22. Kast, W.M., Brandt, R.M.P., Sidney, J., Drijfhout, J.W., Kubo, R.T., Melief, C.J.M. and Sette, A. (1994) Role of HLA-A motifs in identification of potential CTL epitopes in human papillomavirus type 16 E6 and E7 proteins. *J. Immunol.* **152**, 3904-3912.
23. Abrams, S.I., Stanziale, S.F., Lunin, S.D., Zaremba, S. and Schlom, J. (1996) Identification of overlapping epitopes in mutant ras oncogene peptides that activate CD4⁺ and CD8⁺ T cell responses. *Eur. J. Immunol.* **26**, 435-443.
24. Cheever, M.A., Disis, M.L., Bernhard, H., Gralow, J.R., Hand, S.L., Huseby, E.S., Takahashi, M. and Chen, W. (1995) Immunity to oncogenic proteins. *Immunol. Rev.* **145**, 33-59.
25. Uenaka, K., Ono, T., Akisawa, T., Wada, H., Yasuda, T. and Nakayama, E. (1994) Identification of a unique antigen peptide pRL1 on BALB/c RL1 leukemia recognized by cytotoxic T lymphocytes and its relation to the Akt oncogene. *J. Exp. Med.* **180**, 1599-1607.
26. Mandelboim, O., Berke, G., Fridkin, M., Feldman, M., Eisenstein, M. and Eisenbach, L. (1994) CTL induction by a tumor-associated antigen

- octapeptide derived from a murine lung carcinoma. *Nature (Lond.)* **369**, 67-71.
27. Cox, A., Skipper, J., Chen, Y., Henderson, R., Darrow, T., Shabanowitz, J., Engelhard, V., Hunt, D. and Slingluff, C. (1994) Identification of a peptide recognized by five melanoma-specific human cytotoxic T cell lines. *Science* **264**, 716-719.
 28. Blake, J., Johnston, J., Hellstrom, K.E., Marquardt, H. and Chen, L. (1996) Use of combinatorial peptide libraries to construct functional mimics of tumor epitopes recognized by MHC class I restricted T lymphocytes. *J. exp. Med.* **184**, 121-130.
 29. Boon, T., Cerottini, J.-C., Van den Eynde, B., Van der Bruggen, P. and Van Pel, A. (1994) Tumor antigens recognized by T lymphocytes. *Annu. Rev. Immunol.* **12**, 337-365.
 30. Brichard, V., Van Pel, A., Wölfel, C., Wölfel, E., DePlaen, E., Lethe, B., Coulie, P. and Boon, T. (1993) The tyrosinase gene codes for an antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanoma. *J. Exp. Med.* **178**, 489-485.
 31. Kawakami, Y., Eliyahu, S., Delgado, C., Robbins, P., Rivoltini, L., Topalian, S., Miki, T. and Rosenberg, S. (1994) Cloning of the gene coding for a shared human melanoma antigen recognized by autologous T cells infiltrating into tumor. *Proc. Natl. Acad. Sci. USA* **91**, 3515-3519.
 32. Coulie, P., Brichard, V., Van Pel, A., Wölfel, T., Schneider, J., Traversari, C., Mattei, S., DePlaen, E., Lurquin, C., Szikora, J., Renault, J. and Boon, T. (1994) A new gene coding for a differentiation antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas. *J. Exp. Med.* **180**, 1-4.
 33. Wang, R., Robbins, P., Kawakami, Y., Kang, X. and Rosenberg, S. (1995) Identification of a gene encoding a melanoma tumor antigen recognized by HLA-A31-restricted tumor infiltrating lymphocytes. *J. Exp. Med.* **181**, 799-804.
 34. Kawakami, Y., Eliyahu, S., Delgado, C., Robbins, P., Sakaguchi, K., Appella, E., Yannelli, J., Adema, G., Miki, T. and Rosenberg, S. (1994) Identification of a human melanoma antigen recognized by tumor-rejection. *Proc. Natl. Acad. Sci. USA* **91**, 6458-6462.
 35. Marchand, M., Weynants, P., Rankin, E., Arienti, F., Belli, F., Parmiani, G., Cascinelli, N., Bourland, A., Wanwijck, R., Humblet, Y., Canon, J.L., Naeyaert, J.M., Plagne, R., Deraemaeker, R., Knuth, A., Jager, E., Brasseur, F., Herman, J., Coulie, P.G. and Boon, T. (1995) *Int. J. Cancer* **63**, 883-885.

36. Rosenberg, S.A. et al. (1998) Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nature Med.* **4**, 321-327.
37. Nestlel, F.L., Aljagic, S., Gillietl, M., Sun, Y., Grabbe, S., Dummerl, R., Burge, G. and Schadendorf, D. (1998) Vaccination of melanoma patients with peptide or tumor lysate-pulsed dendritic cells. *Nature Med.* **4**, 328-332.
38. Mandelboim, O., Vadai, E., Fridkin, M., Katz-Hillel, A., Feldman, M., Berke, G. and Eisenbach, L. (1995) Regression of established murine carcinoma metastases following vaccination with tumor-associated antigen. *Nature Med.* **1**, 1179-1183.
39. Mayordomo, J.I., Zorina, T., Storkus, W.J., Zitvogel, L., Cellzsi, C, Falo, L.D., Melief, C.J., Ildstadst, W.M., Deleo, A.B.(1995. Bone marrow derived dendritic cells pulsed with synthetic tumor peptides elicit protective and therapeutic anti tumor immunity. *Nature Medicine* **1**, 1297-1302.
40. Fearon, E.R., Pardoll, D.M., Itaya, T., Golumbek, P., Levitsky, H.I., Simons, J.W., Karasuyama, H., Vogelstein, B., and Frost, P. (1990) Interleukin-2 production by tumor cells by passes T helper function in generation of an anti-tumor response. *Cell* **60**, 397-403.
41. Tepper, R.I. and Mule, J. (1994) Experimental and clinical studies of cytokine gene modified tumor cells. *Hum. Gene Ther.* **5**, 1530164.
42. Townsend, S.E. and Allison, J.P. (1993) Tumor rejection after direct costimulation of CD8⁺ T cells by B7-transfected melanoma cells. *Science* **259**, 368-370.
43. Darrow, T., Slingluff, C. and Seigler, H. (1989) The role of HLA class I antigens in recognition of melanoma cells by tumor-specific T lymphocytes. Evidence for shared tumor antigens. *J. Immunol.* **142**, 3329-3335.
44. Hom, S., Topalian, S., Simoni, S., Mancini, M. and Rosenberg, S. (1991) Common expression of melanoma tumor-associated antigens recognized by human tumor-infiltrating lymphocytes: analysis by human lymphocyte antigen restrictin. *J. Immunother.* **10**, 153-164.
45. Anichini, A., Maccalli, C., Mortarini, R., Salvi, S., Mazzocchi, Z., Squarcina, P., Herlyn, M. and Parmiani, G. (1993) Melanoma cells and normal melanocytes share antigens recognized by HLA-A2 restricted cytotoxic T cell clones from melanoma patients. *J. Exp. Med.* **177**, 989-998.
46. Bakker, A., Schreurs, M., deBoer, A., Kawakami, Y., Rosenberg, S., Adema, G. and Figdor, C. (1994) Melanocyte lineage-specific antigen gp100 is recognized by melanoma-derived tumor-infiltrating lymphocytes. *J. Exp. Med.* **179**, 1005-1009.

47. Robbins, P., El-Gamil, M., Kawakami, Y., Stevens, E., Yannelli, J. and Rosenberg, S. (1994) Recognition of tyrosinase by tumor infiltrating lymphocytes from a patient responding to immunotherapy. *Cancer Res.* **54**, 3124-3126.
48. Sette, A., Vitiello, A., Reheman, B., Fowler, P., Nayersina, R., Kast, W.M., Melief, C.J., Oseroff, C., Yuan, L., Ruppert, J. et al. (1994) The relationship between class I binding affinity and immunogenicity of potential cytotoxic T cell epitopes. *J. Immunol.* **153**, 5586-5592.
49. Shirai, M., Arichi, T., Nishioka, M., Nomura, T., Ikeda, K., Kawanishi, K., Engelhard, V.H., Feinstone, S.M. and Berzofsky, J.A. (1995) CTL responses of HLA-A2.1-transgenic mice specific for hepatitis C viral peptides predict epitopes for CTL of humans carrying HLA-A2.1. *J. Immunol.* **154**, 2733-1942.
50. Man, S., Newberg, M.H., Crotzer, V.L., Luckey, C.J., Williams, N.S., Chen, Y., Huczko, E.L., Ridge, J.P. and Engelhard, V.H. (1995) Definition of a human T cell epitope from influenza A non-structural protein 1 using HLA-A2.1 transgenic mice. *Int. Immunol.* **7**, 597-605.
51. Rensing, M.E., Sette, A., Brandt, R.M., Ruppert, J., Wentworth, P.A., Hartman, M., Oseroff, C., Grey, H.M., Melief, C.J. and Kast, W.M. (1995) Human CTL epitopes encoded by human papillomavirus type 16 E6 and E7 identified through *in vivo* and *in vitro* immunogenicity studies of HLA-A*0201-binding peptides. *J. Immunol.* **154**, 5934-5943.
52. Wentworth, P.A., Vitiello, A., Sidney, J., Keogh, E., Chesnut, R.W., Grey, H. and Sette, A. (1996) Differences and similarities in the A2.1-restricted cytotoxic T cell repertoire in humans and human leukocyte antigen-transgenic mice. *Eur. J. Immunol.* **26**, 97-101.
53. Barra, C., Gournier, H., Garcia, Z., Marche, P.N., Jouvin-Marche, E., Briand, P., Fillipi, P. and Lemonnier, F.A. (1993) Abrogation of H-2-restricted CTL responses and efficient recognition of HLA-A3 molecules in DBA/2 HLA/A24 responder mice. *J. Immunol.* **150**, 3681-3689.
54. Vitiello, A., Marchesini, D., Furze, J., Sherman, L.A., Chestnut, R.W. (1991) Analysis of HLA restricted influenza specific cytotoxic T lymphocytes response in transgenic mice carrying a chimeric human-mouse class I major histocompatibility complex. *J. Exp. Med.* **173**, 1007-1015.
55. Pascolo, S., Bervas, N., Ure, J.M., Smith, A.G., Lemonnier, F.A. and Rerarnau, B. (HLA-A2.1 restricted education and cytolytic activity of CD8⁺ T lymphocytes from β 2 microglobulin (β 2m) HLA-A2.1 monochain

- transgenic, H-2D^b, β 2m double knockout mice. J. Exp. Med. **185**, 2043-2051.
56. Messing, E.M. and Catalona, W. Urothelial tumors of the urinary tract. In Campbell's Urology, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th editin, WB, Sounders Co. 1998), 2327-2410.
 57. Ioachim-Velogianni, E., Stavropoulos, N.E., Kitsiou, E., Stefanaki, S. and Agnantis, N.J. (1995) Distribution of CD1A-positive Langerhans cells and lymphocytes subsets in transitional cell carcinoma of the urinary bladder. An immunohistological study on frozen sections. J. Pathol. **155**, 401-406.
 58. Thanhauser, A., Bohle, A., Flad, H.D., Ernst, M., Mattern, T. and Ulmer, A.J. (1993) Induction of bacillus Calmette-Gherin activated killer cells from human peripheral blood mononuclear cells against bladder carcinoma cell lines *in vitro*. Cancer Immunol. Immunother. **37**, 105-111.
 59. Patard, H., Muscatelli-Groux, B., Saint, F., Popov, Z., Maille, P., Abbou, C. and Chopin, D. (1996) Evaluation of local immune response after intravesical bacille Calmette Guerin treatment for superficial bladder cancer. Br. J. Urol. **78**, 709-714.
 60. Housseau, F., Zeliskewski, D., Roy, M. et al. (1997) MHC-dependent cytotoxicity of autologous tumor cells by lymphocytes infiltrating urothelial carcinomas. Int. J. Cancer **71**, 585-594.
 61. Moll, R., Wu, X-R., Lin, J-H. and Sun, T-T. (1995) Uroplakins, specific membrane proteins of urothelial umbrella cells, as histological markers of metastatic transitional cell carcinomas. Am. J. Pathol. **147**, 1383-1397.
 62. Mandelboim, O., Bar-Haim, E., Vadai, E., Fridkin, M. and Eisenbach, L. (1997) Identification of shared Tumor-Associated Antigen peptides between two spontaneous lung carcinomas. J. Immunol. **159**, 6030-6036.
 63. Boring, C.C., Squires, T.S., Tong, T. and Montgomery, S. (1994) Cancer statistics 1994 CA **43**, 7-26.
 64. Carter, H.B., Pearson, J.D., Metter, J.E. et al. (1992) Longitudinal evaluation of prostate specific antigen levels in men with and without prostate disease. JAMA **267**, 2215-2220.
 65. Partin, A.W., Poind, C.R., Clemens, J.Q. et al. (1993) Serum PSA after anatomic radical prostatectomy. Urol. Clin. North Am. **20**, 713-725.
 66. Epstein, J.I., Partin, A.W., Sauvageot, J. and Walsh, P.C. (1996) Prediction of progression following radical prostatectomy. A multivariate analysis of 721 men with long term follow up. Am. J. Surg. Pathol. **20**, 286-292.

67. Eisenberger, M.H. (1998) Chemotherapy for hormone resistant prostate cancer. In Campbell's Urology, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th edition, WB, Saunders Co.) 2645-2658.
68. Schroder, F.H. (1998) Endocrine treatment of prostate cancer. In Campbell's Urology, Eds. Walsh, P.C., Retik, A.B., Vaughan, E.D. and Wein, A.J. (7th edition, WB, Saunders Co.) 2627-2644.
69. Xue, B.H., Zhang, Y., Sosman, J.A. and Peace, D.J. (1997) Induction of human cytotoxic T lymphocytes specific for prostate specific antigen. *The Prostate* **30**, 73-78.
70. Correale, P., Walmsley, K., Nieroda, C., Zhu, M., Schlom, J. and Tsang, K.Y. (1997) *In vitro* generation of human cytotoxic T lymphocytes specific for peptides derived from prostate specific antigen. *J. Natl. Cancer Inst.* **89**, 272-275.
71. Tjoa, B., Bounton, A., Kenny, G., Ragde, H., Misrock, S.L. and Murphy, G. (1996) Presentation of prostate tumor antigens by dendritic cells stimulates T cell proliferation and cytotoxicity. *Prostate* **28**, 65-69.
72. Murphy, G., Tjoa, B., Ragde, H., Kenny, G. and Bounton, A. (1996) Phase I clinical trial: T cell therapy for prostate cancer using autologous dendritic cells pulsed with HLA-A0201 specific peptides from Prostate Specific Membrane Antigen. *The Prostate* **29**, 371-380.
73. Bonandona, G. (1996) Future development in adjuvant systemic therapy for high risk breast cancer. *Cancer Res.* **140**, 227-234.
74. Barnd, D.L., Lan, M., Metzger, R., Finn, O.J. (1989). Specific, MHC-unrestricted recognition of tumor associated mucins by human cytotoxic T cells. *Proc. Natl. Acad. SCI. USA* **86**, 7159-7163.
75. Agrawal, B., Reddish, M., Longenecker, B.M. (1996). *In vitro* induction of MUC-1 peptide specific type 1 T lymphocyte and cytotoxic T lymphocyte responses from healthy multiparous donors. *J. Immunol* **157**, 2089-2095.
76. Agrawal, B., Reddish, M.A., Krantz, M.J., Langenecker, B.M. (1995). Does pregnancy immunize against breast cancer? *Cancer Res.* **55**, 2257-2261.
77. Fisk, B., Blevins, T.L., Taylor-Wharton, J., Ionnides, C.G., (1995). Identification of an immunodominant peptide of HER2/neu protooncogene recognized by ovarian Tumor specific cytotoxic T lymphocyte lines. *J. Exp. Med.* **181**, 2109-2117.
78. Linehan, C., Goedegebuure, P.S., Peoples, G.E., Rogers, S.O., Eberlein, P.J. (1995) Tumor specific and HLA-A2 restricted cytotoxicity by tumor-associated lymphocytes in human metastatic breast cancer. *J. Immunol* **155**, 4486-4491.

79. Larocca, D., Peterson, J.A., Urrea, R., Kuniyoshi, J., Bistrain, A.M. and Ceriani, R.L. (1991) A Mr 46,000 human milk fat globule protein that is highly expressed in human breast tumors contains factor VIII-like domains. *Cancer Res.* **15**, 4994-4998.
80. Taylor, M.R., Couto, J.R., Scallan, C.D., Ceriani, R.L. and Peterson, J.A. (1997) Lactadherin (formerly BA46), a membrane-associated glycoprotein expressed in human milk and breast carcinomas, promotes Arg-Gly-Asp (RGD)-dependent cell adhesion. *DNA Cell Biol.* **16**, 861-869.
81. Ceriani, R.L., Blank, E.W., Couto, J.R., and Peterson, J.A. (1995) Biological activity of two humanized antibodies against two different breast cancer antigens and comparison to their original murine forms. *Cancer Res.* **155**, 5852-S5856S.
82. Couto, J.R., Blank, E.W., Peterson, J.A., Ceriani, and R.L. (1995) Anti-BA46 monoclonal antibody Mc3: humanization using a novel positional consensus and *in vivo* and *in vitro* characterization. *Cancer Res.* **55**, 1717-1722.
83. Peterson, J.A., Couto, J.R., Taylor, M.R., and Ceriani R.L. (1995) Selection of tumor specific epitopes on target antigens for radioimmunotherapy of breast cancer. *Cancer Res.* **55**, (Suppl.) 5847-5851.
84. Graham, G.A., Burchell, J.M., and Taylor-Papadimitriou, J. (1996) The polymorphic epithelial Mucin: potential as an immunogen for a cancer vaccine. *Cancer Immunol. Immunother.* **42**, 71.
85. Ho, S.B., Niehans, G.A., Lyftogt, C., Yan, P.S., Cherwitz, D.L., Gum, E.T., Dahiya, R., and Kim, Y.S. (1993) Heterogeneity of Mucin gene expression in normal and neoplastic tissues. *Cancer Res* **1**, 641-651
86. Walsh, M.D., Hohn, B.G., Thong, W., Devine, P.L., Gardiner, R.A., Samaratunga, M.L. and McGuckin, M.A. (1994) Mucin expression by transitional cell carcinomas of the bladder. *Br. J. Urol.* **73**, 256-62
87. Noto, H., Takahashi, T., Makiguchi, Y., Hayashi, T., Hinoda, Y. and Imai, K. (1997) Cytotoxic T lymphocytes derived from bone marrow mononuclear cells of multiple myeloma patients recognize an underglycosylated form of MUC1 Mucin. *Int Immunol* **9**(5):791-798
88. Spicer, A.P., Rowse, G.J., Linder, T.K. and Gendler, S.J. (1995) Delayed mammary tumor progression in Muc-1 null mice. *J. Biol. Chem.* **270**, 30093.
89. McGuckin, M.A., Walsh, M.D., Hohn, B.G., Ward, B.G. and Wright, R.G. (1995) Prognostic significance of Muc-1 epithelial Mucin expression in breast cancer. *Hum. Pathol.* **26**, 432.

90. Wreschner, D.H., Hareuveni, M., Tsarfaty, I., Smorodinsky, N., Horev, J., Zaretsky, J., Kotkes, P., Weiss, M., Lathe, R. and Dion, A. (1990) Human epithelial tumor antigen sequences. Differential splicing may generate multiple protein forms. *Eur. J. Biochem.* **189**, 463.
91. Tsarfaty, I., Hareuveni, M., Horev, J., Zaretsky, J., Weiss, M., Jeltsch, J.M., Garnier, J.M., Lathe, R., Keydar, I. and Wreschner, D.H. (1990) Isolation and characterization of an expressed hypervariable gene coding for a breast caantigen. *Gene* **93**, 313.
92. Domenech, N. and Finn, O.J. (1995) *In vitro* studies of MHC-restricted recognition of a peptide from the MUC1 tandem repeat domain by CTL. *FASEB J.* **9**, 1023.
93. Barnd, D.I., Lan, M.S., Mezgar, R.S. and Finn, O.J. (1989) Specific, major histocompatibility complex-unrestricted recognition of tumor associated mucins by human cytotoxic T cells. *Proc. Natl. Acad. Sci. USA* **86**, 7159.
94. Domenech, N., Henderson, R.A. and Finn, O.J. (1995) Identification of a HLA-A11-restricted epitope from the tandem repeat domain of the epithelial tumor antigen Mucin. *J Immunol* **15**, 4766.
95. Apostolopoulos, V., Karanikas, V., Haurum, J.S. and McKenzie, I.F. (1997) Induction of HLA-A2-restricted CTLs to the Mucin 1 human breast cancer antigen. *J. Immunol.* **159**, 5211.
96. Mackenzi, P.Y. and Longenecker, B.M. (1991) Specific immunosuppressive activity of epiglycanin, a Mucin-like glycoprotein secreted by a murine mammary adenocarcinoma (TA3-HA). *Cancer Res.* **15**, 1170.
97. Gimmi, C.D., Morrison, B.W., Mainpric, B.A., Gribben, J.G., Boussiotis, V.A., Freeman, G.J., Park, S.Y., Watanabe, M., Gong, J., Hayes, D.F., Kufe, D.W. and Nadler, L.M. (1996) Breast cancer-associated antigen, DF3/MUC1, induces apoptosis of activated human T cells. *Nat. Med. Dec.* **2**, 1367.
98. Agrawal, B., Krantz, M.J., Reddish, MA., Longenecker and B.M. (1998) Cancer-associated MUC1 Mucin inhibits human T-cell proliferation, which is reversible by IL-2. *Nat Med.* **4**, 43.
99. Spooner, R.A., Deonarain, M.P. and Epenetos, A.A., DNA vaccination for cancer treatment. *Gene Therapy* **2**, 173-180, 1995.
100. Ulmer, J.B., Donnelly, J.J., Parker, S.E., Rhodes, G.H., Felgner, P.L., Dwaraki, V.J., Gromkowski, S.H., Deck, R.R., DeWitt, C.M., Friedman, A., Hawe, L.A., Leander, K.R., Martinez, D., Perry, H.C., Shiver, J.W., Montgomery, D.L. and Liu, M.A., Heterologous protection against

- influenza by injection of DNA encoding a viral protein. *Science* 259, 1745-1749, 1993.
101. Conry, R.M., LoBuglio, A.F., Kantor, J., Schlom, J., Loechel, F., Moore, S.E., Sumerel, L.A., Barlow, D.L., Abrams, S. and Curiel, D.T., Immune response to a carcinoembryonic antigen polynucleotide vaccine. *Cancer Res.*, 54, 1164-1168, 1994.
 102. Conry, R.M., LoBuglio, A.F., Loechel, F., Moore, S.E., Sumerel, L.A., Barlow, D.L., Pike, J. and Curiel, D.T. A carcinoembryonic antigen polynucleotide vaccine for human clinical use: *Cancer Gene Therapy* 2, 33-38, 1995.
 103. Acres, R.B., Hareuveni, M., Balloul, J.-M. and Kieny, M.-P. Vaccinia virus MUC1 immunization of mice: immune response and protection against the growth of murine tumors bearing the MUC1 antigen. *J. Immunother.* 14, 136-143, 1993.
 104. Oldstone, M.B., Tishon, A., Eddleston, M., de la Torre, J.C., McKee, T. and Whitton, J.L. Vaccination to prevent persistent viral infection. *J. Virol.* 67, 4372-4378, 1993.
 105. Whitton, J.L., Sheng, N., Oldstone, M.B. and McKee, T.A., A "string-of-beads" vaccine, comprising linked minigenes, confers protection from lethal-dose virus challenge. *J. Virol.* 67, 348-352, 1993.
 106. Thomson, S.A., Khanna, R., Gardner, J., Burrows, S.R., Coupar, B., Moss, D.J. and Suhrbier, A., Minimal epitopes expressed in a recombinant polyepitope protein are processed and presented to CD8+ cytotoxic T cells: implications for vaccine design. *Proc. Natl. Acad. Sci. USA* 92, 5845-5849, 1995.
 107. Thomson, S.A., Elliott, S.L., Sherritt, M.A., Sproat, K.W., Coupar, B.E., Scalzo, A.A., Forbes, C.A., Ladhams, A.M., Mo, X.Y., Tripp, R.A., Doherty, P.C., Moss, D.J. and Suhrbier, A., Recombinant polyepitope vaccines for the delivery of multiple CD8 cytotoxic T cell epitopes. *J. Immunol.* 157, 822-826, 1996.
 108. Thomson, S.A., Sherritt, M.A., Medveczky, J., Elliott, S.L., Moss, D.J., Fernando, G.J., Brown, L.E., and Suhrbier, A., Delivery of multiple CD8 cytotoxic T cell epitopes by DNA vaccination. *J. Immunol.* 160, 1717-1723, 1998.
 109. Boczkowski, D., Nair, S.K., Snyder, D. and Gilboa, E., Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J. Exp. Med.* 184, 465-472, 1996.

110. Mayordomo, J.I., Zorina, T., Storkus, W.J., Zitvogel, L., Celluzzi, C., Faló, L.D., Melief, C.J., Ildstad, S.T., Kast, W.M., Deleo, A.B., et al., Bone marrow-derived dendritic cells pulsed with synthetic tumor peptides elicit protective and therapeutic antitumor immunity. *Nature Med.* 1, 1297-1302, 1995.
111. Mayordomo, J.I., Loftus, D.J., Sakamoto, H., Decesare, C.M., Appasamy, P.M., Lotze, M.T., Sorkus, W.J., Appella, E., Deleo, A.B., and De Cesare, C.M., Therapy of murine tumors with P53 wild-type and mutant sequence peptide-based vaccines. *J. Exp. Med.* 183, 1357-1365, 1996.
112. Mandelboim, O., Berke, G., Fridkin, M., Feldman, M., Eisenstein, M., and Eisenbach, L., CTL induction by a tumor-associated antigen octapeptide derived from a murine lung carcinoma. *Nature* 369, 67-71, 1994.
113. Mandelboim, O., Vadai, E., Fridkin, M., Katz-Hillel, A., Feldman, M., Berke, G., and Eisenbach, L., Regression of established murine carcinoma metastases following vaccination with tumour-associated antigen peptides. *Nat. Med.* 1, 1179-1183, 1995.
114. Mukherji, B., Chakraborty, N.G., Yamasaki, S., Okino, T., Yamase, H., Sporn, J.R., Kurtzman, S.K., Ergin, M.T., Ozols, J., Meehan, J., et al., Induction of antigen-specific cytolytic T cells in situ in human melanoma by immunization with synthetic peptide-pulsed autologous antigen presenting cells. *Proc. Natl. Acad. Sci. USA* 92, 8078-8082, 1995.
115. Gjertsen, M.K., Bakka, A., Breivik, J., Saeterdal, I., Gedde-Dahl, Stokke, K.T., Solheim, B.G., Egge, T.S., Soreide, O., Thorsby, E., and Gaudernack, G., Ex vivo ras peptide vaccination in patients with advanced pancreatic cancer: results of a phase I/II study. *Int. J. Cancer* 65, 450-453, 1996.
116. Murphy, G., Tjoa, B., Ragde, H., Kenny, G. and Boynton, A., Phase I clinical trial: T-cell therapy for prostate cancer using autologous dendritic cells pulsed with HLA-A0201-specific peptide from prostate-specific membrane antigen. *Prostate* 29, 371-380, 1996.
117. Parker, K.C., Bednarek, M.A., Coligan, J.E., Scheme for ranking potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J Immunol.* 152, 163-175, 1994.

SEQUENCE LISTING

(2) INFORMATION FOR SEQ ID NO:1:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
 Phe Leu Leu Val Leu Gly Phe Ile Ile
 5 9

(2) INFORMATION FOR SEQ ID NO:2:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:
 Val Leu Pro Ser Val Ala Met Phe Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:3:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
 Leu Val Leu Gly Phe Ile Ile Ala Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:4:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
 Lys Val Val Thr Ser Ser Phe Val Val
 5 9

(2) INFORMATION FOR SEQ ID NO:5:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
 Leu Val Pro Gly Thr Lys Phe Tyr Ile
 5 9

(2) INFORMATION FOR SEQ ID NO:6:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
 Leu Leu Pro Ile Arg Thr Leu Pro Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
 Tyr Leu Val Lys Lys Gly Thr Ala Thr
 5 9

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
 Ser Leu Phe Ala Glu Thr Ile Trp Val
 5 9

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
 Met Leu Ile Ala Met Tyr Phe Tyr Thr
 5 9

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:
 Leu Met Trp Thr Leu Pro Val Met Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:
 Met Leu Ile Val Tyr Ile Phe Glu Cys
 5 9

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid

(C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:
 Tyr Ile Phe Glu Cys Ala Ser Cys Ile
 5 9

(2) INFORMATION FOR SEQ ID NO:13:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:
 Leu Val Leu Met Leu Ile Val Tyr Ile
 5 9

(2) INFORMATION FOR SEQ ID NO:14:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:
 Ala Leu Cys Arg Arg Ser Met Val
 5 9

(2) INFORMATION FOR SEQ ID NO:15:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:
 Leu Leu Ser Gly Leu Ser Leu Phe Ala
 5 9

(2) INFORMATION FOR SEQ ID NO:16:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:
 Phe Leu Leu Val Val Gly Leu Ile Val
 5 9

(2) INFORMATION FOR SEQ ID NO:17:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:
 Leu Val Val Gly Leu Ile Val Ala Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:18:
 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:
 Lys Val Val Lys Ser Asp Phe Val Val
 5 9

(2) INFORMATION FOR SEQ ID NO:19:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
 Thr Leu Pro Val Gln Thr Leu Pro Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:20:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
 Asp Leu His Val Ile Ser Asn Asp Val
 5 9

(2) INFORMATION FOR SEQ ID NO:21:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
 Val Leu Val His Pro Gln Trp Val Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:22:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
 Phe Leu Arg Pro Gly Asp Asp Ser Ser
 5 9

(2) INFORMATION FOR SEQ ID NO:23:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:
 Ala Leu Gly Thr Thr Cys Tyr Ala Ser
 5 9

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Lys Leu Gln Cys Val Asp Leu His Val

5

9

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Glu Leu Ala His Tyr Asp Val Leu Leu

5

9

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Asn Leu Asn Gly Ala Gly Asp Pro Leu

5

9

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Thr Leu Arg Val Asp Cys Thr Pro Leu

5

9

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Met Met Asn Asp Gln Leu Met Phe Leu

5

9

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Ala Leu Phe Asp Ile Glu Ser Lys Val

5

9

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Leu Leu His Glu Thr Asp Ser Ala Val
 5 9

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Val Leu Ala Lys Glu Leu Lys Phe Val
 5 9

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Ile Leu Leu Trp Gln Pro Ile Pro Val
 5 9

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Asp Leu Phe Gly Ile Trp Ser Lys Val
 5 9

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Pro Leu Glu Arg Phe Ala Glu Leu Val
 5 9

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:
 Lys Gln Gly Asn Phe Asn Ala Trp Val
 5 9

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:
 Asn Leu Leu Arg Arg Met Trp Val Thr
 5 9

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:
 Asn Leu Phe Glu Thr Pro Ile Leu Ala
 5 9

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:
 Asn Leu Phe Glu Thr Pro Val Glu Ala
 5 9

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:
 Gly Leu Gln His Trp Val Pro Glu Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:
 Val Gln Phe Val Ala Ser Tyr Lys Val
 5 9

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid

(C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:
 Arg Leu Leu Ala Ala Leu Cys Gly Ala
 5 9

(2) INFORMATION FOR SEQ ID NO:42:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:
 Leu Leu Leu Leu Thr Val Leu Thr Val
 5 9

(2) INFORMATION FOR SEQ ID NO:43:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:
 Leu Leu Leu Thr Val Leu Thr Val Val
 5 9

(2) INFORMATION FOR SEQ ID NO:44:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:
 Phe Leu Ser Phe His Ile Ser Asn Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:45:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:
 Leu Leu Val Leu Val Cys Val Leu Val
 5 9

(2) INFORMATION FOR SEQ ID NO:46:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:
 Ala Leu Leu Val Leu Val Cys Val Leu
 5 9

(2) INFORMATION FOR SEQ ID NO:47:
 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:
 Ser Leu Ser Tyr Thr Asn Pro Ala Val
 5 9

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:
 Asn Leu Thr Ile Ser Asp Val Ser Val
 5 9

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:
 Ala Leu Ala Ser Thr Ala Pro Pro Val
 5 9

WHAT IS CLAIMED IS:

1. A vaccine composition comprising at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).
2. The vaccine composition of claim 1, wherein said Uroplakin is selected from the group consisting of Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.
3. The vaccine composition of claim 1, wherein said at least one tumor associated antigen peptide derived from said Mucin is from a non tandem repeat array of said Mucin.
4. The vaccine composition of claim 1, wherein said at least one tumor associated antigen peptide derived from said Mucin is from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of said Mucin.
5. The vaccine composition of claim 1, wherein said Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1) are each independently of a mammal.
6. The vaccine composition of claim 5, wherein said mammal is selected from the group consisting of humanoid and rodent.
7. The vaccine composition of claim 1, wherein each of said at least one tumor associated antigen peptide includes a number of residues including 2-3 residues which form a primary anchor for interaction with MHC class I molecules and 2-3 residues which bind to T cell receptors.
8. The vaccine composition of claim 7, wherein said number is selected from the group consisting of 7, 8, 9, 10 and 11.

9. The vaccine composition of claim 1, further comprising a carrier, wherein said at least one tumor associated antigen peptide is presented in context of said carrier.

10. The vaccine composition of claim 9, wherein said carrier is selected from the group consisting of a proteinaceous carrier to which said at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.

11. The vaccine composition of claim 1, wherein at least one of said tumor associated antigen peptides includes at least one non-natural modification rendering peptides more stable in a body.

12. The vaccine composition of claim 1, wherein said at least one non-natural modification rendering peptides more stable in said body is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

13. The vaccine composition of claim 1, wherein at least one of said tumor associated antigen peptides includes at least one non-natural modification rendering peptides more immunogenic.

14. The vaccine composition of claim 1, wherein said at least one non-natural modification rendering peptides more immunogenic is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

15. The vaccine composition of claim 1, wherein said at least one tumor associated antigen peptide is selected from SEQ ID NOs: 1-49 and effective homologues and analogs thereof.

16. The vaccine composition of claim 1, wherein the composition is effective in prevention or cure of cancer or cancer metastases.

17. The vaccine composition of claim 16, wherein said cancer is a carcinoma.

18. The vaccine composition of claim 16, wherein said cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, melanoma, colon, stomach and head and neck cancer.

19. A method of vaccination for prevention or cure of cancer comprising the steps of administering a composition including at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

20. The method of claim 19, wherein said Uroplakin is selected from the group consisting of Uroplakin II, Uroplakin Ia, Uroplakin III and Uroplakin Ib.

21. The method of claim 19, wherein said at least one tumor associated antigen peptide derived from said Mucin is from a non tandem repeat array of said Mucin.

22. The method of claim 19, wherein said at least one tumor associated antigen peptide derived from said Mucin is from a region selected from the group consisting of a signal peptide, a cytoplasmic domain and an extracellular domain of said Mucin.

23. The method of claim 19, wherein said Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1) are each independently of a mammal.

24. The method of claim 23, wherein said mammal is selected from the group consisting of humanoid and rodent.

25. The method of claim 19, wherein each of said at least one tumor associated antigen peptide includes a number of residues including 2-3 residues which form a primary anchor for interaction with MHC class I molecules and 2-3 residues which bind to T cell receptors.

26. The method of claim 25, wherein said number is selected from the group consisting of 7, 8, 9, 10 and 11.

27. The method of claim 19, further comprising a carrier, wherein said at least one tumor associated antigen peptide is presented in context of said carrier.

28. The method of claim 27, wherein said carrier is selected from the group consisting of a proteinaceous carrier to which said at least one tumor associated antigen peptide is linked, an adjuvant, a protein or a recombinant protein and an antigen presenting cell.

29. The method of claim 19, wherein at least one of said tumor associated antigen peptides includes at least one non-natural modification rendering peptides more stable in a body.

30. The method of claim 19, wherein said at least one non-natural modification rendering peptides more stable in said body is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

31. The method of claim 19, wherein at least one of said tumor associated antigen peptides includes at least one non-natural modification rendering peptides more immunogenic.

32. The method of claim 19, wherein said at least one non-natural modification rendering peptides more immunogenic is selected from the group consisting of peptoid modification, semipeptoid modification, cyclic peptide modification, N terminus modification, C terminus modification, peptide bond modification, backbone modification and residue modification.

33. The method of claim 19, wherein said at least one tumor associated antigen peptide is selected from SEQ ID NOs: 1-49 and effective homologues and analogs thereof.

34. The method of claim 19, wherein said cancer is a carcinoma.

35. The method of claim 19, wherein said cancer is selected from the group consisting of breast, bladder, prostate, pancreas, ovary, thyroid, melanoma, colon, stomach and head and neck cancer.

36. A DNA vaccine composition comprising at least one polynucleotide encoding a tumor associated antigen peptide derived from a sequence encoding a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

37. The DNA vaccine composition of claim 36, wherein said at least one polynucleotide is a part of a longer polynucleotide designed to encode a fused protein product from which said tumor associated antigen peptide is cleavable by a protease.

38. A cellular vaccine composition comprising an antigen presenting cell presenting at least one tumor associated antigen peptide derived from a protein selected from the group consisting of Uroplakin (UP), Prostate specific antigen (PSA), Prostate specific membrane antigen (PSMA), Prostate acid phosphatase (PAP), Lactadherin (BA46) and Mucin (MUC1).

39. The cellular vaccine composition of claim 38, wherein said antigen presenting cell is selected from the group consisting of a dendritic cell, a macrophage, a B cell and a fibroblast.

40. The cellular vaccine composition of claim 38, wherein presenting said at least one tumor associated antigen peptide is effected by a method selected from the group consisting of:


- (a) transducing said antigen presenting cell with at least one polynucleotide encoding said at least one tumor associated antigen peptide;
- (b) loading said antigen presenting cell with at least one polynucleotide encoding said at least one tumor associated antigen peptide;
- (c) loading said antigen presenting cell with said at least one tumor associated antigen peptide;


- (d) loading said antigen presenting cell with at least one longer polypeptide including said at least one tumor associated antigen peptide



Mark M. Friedman
Advocate, Patent Attorney
Haomanim 7 street
Beit Samueloff, 1st floor
67897 Tel Aviv

Fig. 1









Fig. 3

HhD mice immunized with Uroplakin II peptides



Targets

HhD mice immunized with TCC peptide extract




Fig. 6




Fig. 7





Fig 8

HhD mice immunized with PAP peptides



F 9

HhD mice immunized with PSA peptides




Fig. 10

HhD transgenic mice immunized with PSMA peptides




Fig. 11

HhD mice immunized with HhD transfected, DU 145 cells




Fig. 12

HhD mice immunized with prostate carcinoma peptide extract




Fig. 13


BA46 peptides binding analysis

Fig. 11
BA46 peptides immunogenicity




Fig. 15

Peptide specific CTL responses in breast tumor extract-immunized mice




Fig. 16

Breast tumor extract versus breast normal tissue CTL responses
in breast tumor extract and BA-46 peptide-immunized mice




Fig. 17 HLA-A2.1 (HHD) specific CTL responses in breast tumor extract and BA-46 peptide-immunized mice




Fig. 18


MUC-1 peptides binding analysis

Fig. 19

Breast Tumor extract versus breast Normal tissue CTL responses in MUC-1 peptide immunized mice.





Fig. 20

HLA-A2.1 (HHD) specific CTL responses in breast
tumor extract and MUC-1 peptide-immunized mice






Fig. 21




Fig. 22